

**INDOXACARB****TO CAUSE A HAZARD IN THE USE STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE****PRECAUTIONS:**

- 1 Handle with care. Avoid splashing/contact. Harmful if swallowed or inhaled , mild irritant to eyes and skin.
- 2 Wear full protective clothing, i.e., rubber gloves, overalls, hood, eye-glasses and respirator when handling this product.
- 3 **DO NOT** eat drink or smoke while mixing or applying this insecticide.
- 4 Remove protective clothing on completion of spraying and wash face and hands thoroughly with soap and water.
- 5 Dangerous to wild animals, fish, bees, birds and other animals.
- 6 **DO NOT** contaminate drinking pools, dams, rivers and other waterways.
- 7 Dangerous to livestock and domestic pets.
- 8 **KEEP OUT OF REACH OF CHILDREN , ANIMALS , AND UNINFORMED PERSONS ..**
- 9 **KEEP APART FROM FOOD AND FOODSTUFFS, DRINKING WATER AND EATING UTENSILS.**
- 10 Store in original container with lid tightly closed and **KEEP UNDER LOCK AND KEY.**
- 11 Destroy the empty container by perforation and flattening after rinsing the container at least three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank ..
- 12 .Clean applicator thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers or dams.

**WARNING:**

Allow the following withholding period between last application and grazing/harvesting:

|                                   |         |
|-----------------------------------|---------|
| Tomatoes                          | 3 days  |
| Apples , Pears, Plums and Peaches | 14 days |

**SYMPTOMS OF POISONING:**

Should accidental poisoning occur, perform first aid and take the patient immediately to the nearest physician and show this label. Following ingestion nausea, vomiting confusion and lethargy may occur. There is no specific antidote.

**FIRST AID:** If inhaled , remove patient to a well ventilated area .. If ingested **DO NOT** induce vomiting and get medical attention. If skin is contaminated: wash thoroughly with soap and cold water and remove contaminated clothing.  
If eyes are contaminated: flush with clean water for 15 minutes.

**WARRANTY:**

This product and recommendations concerning its use should achieve excellent results, provided the instructions on this label are followed in full. However, since the performance of crop chemicals can be influenced by factors beyond the control of the manufacturer and seller no warranties are given with regard to the use of this product. The manufacturer and seller accept no responsibility for damage or injury of loss arising from the handling, use or storage of this product.

**ACCLIMA 240 EC**

Insecticide

Reg.No: 21-D-166-  
104-D-173-26

Contents  
1 litre

**POISON****COMPOSITION****Mass/vol**

|                   |            |
|-------------------|------------|
| Indoxacarb        | 90 g/      |
| acetamiprid       | 150g/l l   |
| Inert ingredients | to 1 litre |

CHEMICAL GROUP: Oxadiazine  
Acetamidine

***A stomach , contact  
and systemic  
insecticide for the  
control of various  
insect pests on crops  
as indicated.***

REGISTERED AND DISTRIBUTED BY:  
Refiloe Trading (Pvt) Ltd  
P O Box AY 201, AMBY  
Bayer Complex  
1 Martin Road, Msasa  
Harare, Zimbabwe  
Tel: +263-4-446882/6

Date of Manufacture: See container  
Batch No: See container  
Expiry Date: 2 years from date of manufacture

**DIRECTIONS FOR USE****COMPATIBILITY;**

Compatibility of ACETAMIPRID/INDOXACARB may be influenced by several factors and a physical compatibility test must be performed before spraying . .

**MIXING INSTRUCTIONS**

Fill the tank 25-50% full with water and add the required amount of ACETAMIPRID/INDOXACARB to the tank while it is filled with water. Continue agitating the spray mixture constantly during mixing and spraying operations.

| CROP and PEST                                                                                                                    | DOSAGE            | APPLICATION DIRECTIONS                                                                                                                                                                                                                                         |
|----------------------------------------------------------------------------------------------------------------------------------|-------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>APPLES</b><br>Bollworm larva                                                                                                  | 40ml/100 lt water | Apply as a full cover spray before Larva enter fruit .Repeat 10-14 days later dependent on re-infestation .                                                                                                                                                    |
| Codling moth Larva                                                                                                               | 40ml/100 lt water | Apply as a full cover spray at 75% petal fall at the onset of the first generation of the pest . Apply in a programme of 2 applications with a14 day interval .<br>Do not exceed 4 applications per season .                                                   |
| <b>MAIZE</b><br>Fall Armyworm<br>African bollworm<br>Maize stalk borer                                                           | 400ml/ha          | 400-600L water / ha. Apply twice with a 7 day interval                                                                                                                                                                                                         |
| <b>TOMATOES</b><br>Bollworm larva<br>Tomato semi-looper<br><i>Tuta absoluta</i><br>Aphis                                         | 40ml/100l water   | Apply as a full cover spray at 500-1500 lt spray mix per Ha and spray both sides of trellised tomatoes.<br><br>Commence spraying when first larva appear and follow-up 7 days later , then switch to a registered product with a different mode of action .    |
| <b>POTATOES</b><br>Potato tuber moth<br>Aphis<br>Leafmine r                                                                      | 400ml/ha          | Apply as a full cover spray at 400-600 lt spray mix per Ha<br>Commence when first symptoms appear ( mines or moths) Use in a programme spraying @ 10-14 day intervals dependent on re-infestation .<br>Do not exceed 3 applications of this product per season |
| <b>TOBACCO</b><br>Bollworm larva<br>TRB no .21-24-B-47                                                                           | 400 ml/ha         | Apply as a full cover spray when first larvae appear                                                                                                                                                                                                           |
| <b>BERRIES</b><br>Bollworm larva                                                                                                 | 40ml/100 l water  | Apply as a full cover spray at 500-1000 lt spray mix per Ha                                                                                                                                                                                                    |
| <b>CUCURBITS</b><br>(Pumpkins, Squash,<br>Watermelons, Melons,<br>Baby Marrows, Patty<br>Pans and Cucumbers)<br>African Bollworm | 400ml/ha          | Apply two applications with 7 – 10 day spray intervals when larvae are first noticed.<br><br>Do not exceed two applications of Acclima 250 EC per season.                                                                                                      |
| <b>CRUCIFERAE:</b><br>(Cabbage,<br>Cauliflower,<br>Brussels sprouts,<br>Broccoli)<br>Diamondback moth<br>Aphis                   | 40ml/100 l water  | Apply three times at the interval of 7 days apart in a spray mixture of 500l/ha                                                                                                                                                                                |
| <b>SOYABEAN</b><br>African bollworm<br>Fall Army Worm                                                                            | 400ml/ha          | Apply 250L water / ha. Apply twice with a 7 day interval                                                                                                                                                                                                       |