

TO CAUSE A HAZARD IN THE USE, STORAGE OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE

PRECAUTIONS:

Aro-CungFu should not be applied in a spray solution having a pH of less than 6.5 as phytotoxicity may occur. Avoid inhalation of spray mist and contact with eyes. Wear rubber gloves and face shield when preparing spray mixture. Avoid spray-drift onto adjacent crops, grazing, rivers, dams and areas not under treatment. Do not eat, drink or smoke whilst applying or mixing or before washing hands and face. Prevent contamination of food, feed, drinking water and eating utensils. Clean applicator after use and dispose of wash water where it will not contaminate crops, grazing, rivers and dams. Rinse empty container three times with a volume of water equal to at least one tenth of that of the container and add rinsing to the contents of the spray tank before disposing of the container. Destroy empty container and do not use for any other purpose.

SYMPTOMS OF HUMAN POISONING

Excessive exposure may cause cough, mucous production, shortness of breath, reflecting metal fume fever. If ingested in large quantities symptoms may include nausea and vomiting, abdominal pain and central nervous system depression.

FIRST AID

Inhalation: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention immediately.

Eye contact: Hold eyelids open and flush with water until no evidence of chemical remains (at least 15 – 20 minutes). Get medical attention.

Skin contact: Remove contaminated clothing and shoes. Wash with plenty of soap and water until no evidence of chemical remains (approximately 15 – 20 minutes). Get medical attention.

Ingestion: Drink promptly a large quantity of milk, egg white, gelatine solution or if these are not available, large quantities of water.

NOTE TO PHYSICIAN

There is no specific antidote. Treat symptomatically.

WARNINGS:

If the crop to be treated is intended for export, consult the relevant exporting organization regarding withholding periods.

In the case of crops grown for local consumption allow the following withholding periods:

Table grapes: Do not apply after berries have reached the pea size stage (4–6 weeks prior to harvesting). **Avocados, citrus, Coffee, lettuce, Mangoes, potatoes, Peaches & plums** Allow 14 days between last application and harvesting. **Onions** Do not apply later than 3 days before harvesting. **Tomatoes, cruciferae & beans** Do not apply later than 3 days before harvesting. **Wine grapes** Do not apply later than 21 days before harvesting.

Withholding periods are based on good agricultural practice and when followed, local South African MRL's will be met if the recommended dosage is applied. For fruit destined for export, consult your export agent or supplier for the latest withholding periods relevant to your export market.


AGRO-CUNGFU

Fungicide and Bactericide

Reg. No:

Net Contents:

2



COMPOSITION	%w/w
Copper Hydroxide	37.3 %w/w
Copper Equivalent	24.3 %w/w

CHEMICAL GROUP: Inorganic compound

A suspension concentrate contact fungi- and bactericide to control the listed diseases on the crops as listed.

Registered and Distributed by:

Optimum Agro (Pvt) Ltd
12 Highfield Road,
Southerton, Harare
Zimbabwe.
Tel: 0242774725



Manufactured by:

AECI Plant Health
(a division of AECI Ltd.)
14 Field Road, Lilianton,
Boksburg, 1459, South Africa
Tel +27 11 823 8000
www.aeciph.com



Date of Manufacture: See container

Batch No: See container

Expiry Date: See container

Shelf life: Two years from the date of manufacture if kept in the original unopened container under constant cool and dry conditions.

Harmful when swallowed and fatal if inhaled.

Very toxic to aquatic organisms.

Keep out of reach of children, animals and uninformed persons.

Store away from food and feedstuffs.

Re-entry: Do not enter treated area within 1 day after treatment unless wearing protective clothing.

In case of poisoning – CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM/HER.

ALTHOUGH THIS ADJUVANT HAS BEEN EXTENSIVELY TESTED UNDER A WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY QUALITY OF DILUTION WATER, COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AS WELL AS BY THE METHOD, TIME AND DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

RESISTANCE WARNING:

For resistance management, **Agro-CungFu** is a group code M1 fungicide. Any fungus may contain individuals naturally resistant to **Agro-CungFu** and other group code M1 fungicides. The resistant individuals can eventually dominate the fungus if these fungicides are used repeatedly. The resistant fungi may not be controlled by **Agro-CungFu** or any other group code M1 fungicide.

To delay fungicide resistance: Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with registered products from different fungicide group codes. Integrate other control methods (chemical, cultural, biological) into disease control programmes.

AERIAL APPLICATION:

Aerial application may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 10118 (Aerial Application of Agricultural Remedies). It is important to ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria are met:

Equipment:

Use suitable atomising equipment (hydraulic nozzles or rotary atomisers) that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product through drift.

The operator must use a setup that will produce a droplet spectrum with the lowest possible relative span.

All nozzles / atomisers should be positioned within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.

PXXXX ZimOpt01/24 V1