


AGRO-NIXELA



Read the label carefully before opening.

GROUP 2 FERTILIZER

Reg. No. 388/23

A liquid nutrient complex containing salicylic acid

COMPOSITION:

Calcium (Ca) 2.64 %m/m (3.3 %m/v)
Magnesium (Mg) 0.84 %m/m (1.1 %m/v)
Soluble Potash (K₂O) 5.40 %m/m (6.8 %m/v)
Boron (B) 0.24 %m/m (0.3 %m/v)

Batch No:

Manufacturing date:

Expiry date:

NET VOLUME:

Registered and Distributed by:

Optimum Agro (Pvt) Ltd
12 Highfield Road,
Southerton, Harare
Zimbabwe.
Tel: 0242774725



Manufactured by:

AECI Plant Health
(a division of AECI Ltd.)
14 Field Road, Lilianton,
Boksburg, 1459, South Africa
Tel +27 11 823 8000
www.aeciph.com

For further information see attached pamphlet

UN No.: Not regulated

AGRO-NIXELA

A liquid nutrient complex containing salicylic acid

COMPOSITION:

Calcium (Ca)	2.64 %m/m (3.3 %m/v)
Magnesium (Mg)	0.84 %m/m (1.1 %m/v)
Soluble Potash (K ₂ O)	5.40 %m/m (6.8 %m/v)
Boron (B)	0.24 %m/m (0.3 %m/v)

DIRECTIONS FOR USE: USE ONLY AS INDICATED

AGRO-NIXELA should not be added with any sulphur or phosphite based products

APPLICATION TABLE:

CROP	APPLICATION RATE	APPLICATION METHOD
APPLES, PEARS	200-300 mL/100 ℓ water to a maximum of 3 ℓ/ha	Apply 3-4 applications from 50% flowering and repeat with 21 day intervals. For use on Pears: Do not combine AGRO-NIXELA with other foliar feeds or adjuvants.
CITRUS	200-300 mL/100 ℓ water to a maximum of 3 ℓ/ha	Apply 3 – 4 applications from 100 % petal drop and repeat with 28 day intervals.
CUCURBITS: PUMPKINS, WATERMELONS, MELONS AND CUCUMBERS	500 mL /100 ℓ water to a maximum of 4 ℓ/ha	Apply at flowering and repeat with 21 day intervals.
LETTUCE, CAULIFLOWER, CABBAGE, CHINESE CABBAGE	500 mL/100 ℓ water to a maximum of 3 ℓ/ha	Begin applications prior to onset of head formation and repeat at 14 to 21 day intervals.
PEACHES, NECTARINES, PLUMS, APRICOTS, ALMONDS, OLIVES	200-300 mL/100 ℓ water to a maximum of 3 ℓ/ha	Apply 3-4 applications from 75% petal drop and repeat with 21 to 28 day intervals. If weather conditions favour bacterial infections, additional applications can be considered.
POTATOES	500 mL/100 ℓ water to a maximum of 4 ℓ/ha	Apply at tuber initiation and repeat every 21 to 28 days.
TOMATOES, PEPPERS, BEANS, PEAS, ONIONS	500 mL/100 ℓ water to a maximum of 4 ℓ/ha	Begin applications prior to fruit set and repeat at 14 to 21 day intervals.
TOBACCO	500 mL/100 ℓ water to a maximum of 4 ℓ/ha	Apply 2 – 3 applications bi-weekly, starting at 30 cm height.

IMPORTANT: BEFORE APPLYING AGRO-NIXELA, CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURAL ADVISER ON THE USE THEREOF.