

AGRO-ZNO

Reg. No. 393/23

A water soluble zinc to be used for rapid correction of zinc deficiencies in field - and vegetable crops, subtropical (pecan and avocado) and citrus.

COMPOSITION:

Zinc (Zn).....11.0 %m/m (14.6 %m/v)

Nitrogen (N).....4.8 %m/m (6.4 %m/v)

Registered and Distributed by:

Optimum Agro (Pvt) Ltd
12 Highfield Road,
Southerton, Harare
Zimbabwe.
Tel: 0242774725

For further information see attached pamphlet

UN No.: 3264 - Corrosive liquid, acidic inorganic



Read the label carefully before opening.

GROUP 2 FERTILIZER

Batch No:

Manufacturing date:

Expiry date:

NET VOLUME:



Manufactured by:

AECI Plant Health
(a division of AECI Ltd.)
14 Field Road, Lilianton,
Boksburg, 1459, South Africa
Tel +27 11 823 8000
www.aeciph.com

AGRO-ZNO

A water soluble zinc to be used for rapid correction of zinc deficiencies in field - and vegetable crops, subtropical (pecan and avocado) and citrus.

COMPOSITION:

Zinc (Zn).....11.0 %m/m (14.6 %m/v)

Nitrogen (N).....4.8 %m/m (6.4 %m/v)

Zinc deficiencies occur mainly in alkaline or highly leached acid and sandy soils, following heavy lime or phosphate applications and in soils with an extremely high P-status.

COMPATIBILITY:

AGRO-ZNO should not be combined with pesticides. To improve the efficiency of treatment use, a suitable acidifier-wetter-penetrant such as AGRO-BUFF 5* (refer to AGRO-BUFF 5 label). Mix AGRO-BUFF 5 thoroughly with the water before adding AGRO-ZNO.

RECOMMENDATIONS:

1. Subtropical fruit orchards (pecan, avocado, citrus and others).

Apply as a spring treatment to young foliage (spring flush). In acute cases of deficiency repeated treatments during the summer months may be necessary.

Use 50 - 75ml AGRO-ZNO in 100 ℓ water with a spray volume of 4 000 - 5 000 ℓ water/ha. (2 - 3,75 ℓ AGRO-ZNO/ha). Apply as a cover spray and use higher rate where severe deficiencies occur.

2. Field and vegetable crops (sweet corn, maize, small grain, beans, onions and others).

Apply during early growth stages on young leaves. Under conditions favouring zinc deficiencies successive applications are advisable.

Use 250 - 500 ℓ water/ha and do not exceed 1 ℓ AGRO-ZNO/ha per treatment.

3. Fertigation (tomatoes, lettuce and others).

Where irrigation systems are used for feeding purposes, apply AGRO-ZNO at a rate of 50 - 75ml / 100 ℓ water, with not more than 1 ℓ AGRO-ZNO/ha per treatment. Apply AGRO-ZNO during the early part of the irrigation cycle. Repeated applications may be necessary.

AGRO-BUFF 5: (Contains wetting-spreading agents, acid, pH indicator).