

# Alexin®

## Inorganic Foliar Fertilizer

A liquid nutrient complex containing salicylic acid.

### COMPOSITION:

|           |      |            |            |
|-----------|------|------------|------------|
| Calcium   | (Ca) | 26.4 g/kg  | (33.3 g/l) |
| Magnesium | (Mg) | 8.4 g/kg   | (10.6 g/l) |
| Potassium | (K)  | 45 g/kg    | (56.7 g/l) |
| Boron     | (B)  | 2.375 g/kg | (3.0 g/l)  |

pH: 5 - 8

### DIRECTIONS FOR SAFE DISPOSAL:

Invert empty container over the spray tank and allow draining for at least 30 sec after the flow has slowed down to a drip. Thereafter rinse empty container 3 times with a volume of water equal to at least 10% of that of the container and add the rinses to the contents of the spray tank before disposing of the container. Destroy empty container and do not use for any other purpose.

### SUPPLIER:

**NULANDIS®**, a division of AECI Limited

14 Field Road

Lilianton, Boksburg, 1459, South-Africa

Tel: +27 11 823 8000

www.nulandis.com



A business of  
**AECI**  
Plant & Animal  
Health

### DISTRIBUTOR:

**LACHLAN**

Sunflag Industrial Park, Mombasa road

P.O Box 49470, 00100 Nairobi Kenya

Telephone +254 20 207 3912/3/4

www.lachlanafrica.com



**ALEXIN®** is a registered trademark of Nulandis® (A division of AECI Limited)

Country of origin: South-Africa

### Net Volume:

UN No. Not regulated

Batch No:


Manufacturing date:

Expiry Date:

### STORAGE INSTRUCTIONS:

Store product in a segregated and approved area.

Keep container in a cool, well-ventilated area. Keep container tightly closed and sealed until ready for use.



P7046

# ALEXIN®

A liquid nutrient complex containing salicylic acid.

## COMPOSITION:

|           |      |            |            |
|-----------|------|------------|------------|
| Calcium   | (Ca) | 26.4 g/kg  | (33.3 g/l) |
| Magnesium | (Mg) | 8.4 g/kg   | (10.6 g/l) |
| Potassium | (K)  | 45 g/kg    | (56.7 g/l) |
| Boron     | (B)  | 2.375 g/kg | (3.0 g/l)  |

## Manufacturer & Supplier:

Nulandis, a division of AECI LTD  
14 Field Road, Lilianton, Boksburg, 1459,  
South-Africa  
Tel: +27 11 823 8000



## DISTRIBUTOR:

### LACHLAN

Sunflag Industrial Park, Mombasa road  
P.O Box 49470, 00100 Nairobi Kenya  
Telephone +254 20 207 3912/3/4  
www.lachlanafrika.com



**ALEXIN®** is a registered trademark of AECI Limited (Nulandis®, a division of AECI Ltd)

## DIRECTIONS FOR USE: USE ONLY AS INDICATED

- Alexin should not be added with any sulphur or phosphite based products

## APPLICATION TABLE:

| CROP                                                    | APPLICATION RATE                              | APPLICATION METHOD                                                                                                                                                               |
|---------------------------------------------------------|-----------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| APPLES, PEARS                                           | 200-300 ml/100 l water to a maximum of 3 l/ha | Apply 3-4 applications from 50% flowering and repeat with 21 day intervals. <b>For use on Pears: Do not combine Alexin with other foliar feeds or adjuvants.</b>                 |
| CITRUS                                                  | 200-300 ml/100 l water to a maximum of 3 l/ha | Apply 3 – 4 applications from 100 % petal drop and repeat with 28 day intervals.                                                                                                 |
| CUCURBITS (PUMPKINS, WATERMELONS, MELONS AND CUCUMBERS) | 500 ml /100 l water to a maximum of 4 l/ha    | Apply at flowering and repeat with 21 day intervals.                                                                                                                             |
| LETTUCE, CAULIFLOWER, CABBAGE, CHINESE CABBAGE          | 500 ml/100 l water to a maximum of 3 l/ha     | Begin applications prior to onset of head formation and repeat at 14 to 21 day intervals.                                                                                        |
| PEACHES, NECTARINES, PLUMS, APRICOTS, ALMONDS, OLIVES   | 200-300 ml/100 l water to a maximum of 3 l/ha | Apply 3-4 applications from 75% petal drop and repeat with 21 to 28 day intervals. If weather conditions favour bacterial infections, additional applications can be considered. |
| POTATOES                                                | 500 ml/100 l water to a maximum of 4 l/ha     | Apply at tuber initiation and repeat every 21 to 28 days.                                                                                                                        |
| TOMATOES, PEPPERS, BEANS, PEAS, ONIONS                  | 500 ml/100 l water to a maximum of 4 l/ha     | Begin applications prior to fruit set and repeat at 14 to 21 day intervals.                                                                                                      |
| TOBACCO                                                 | 500 ml/100 l water to a maximum of 4 l/ha     | Apply 2 – 3 applications bi-weekly, starting at 30 cm height.                                                                                                                    |

**IMPORTANT: BEFORE APPLYING ALEXIN, CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURAL ADVISER ON THE USE THEREOF.**