

Alexin®



FPA Registration No.: **1-1IF-8348**

A liquid nutrient complex with the following content:

Potassium (K₂O)..... 6.9%
Calcium (CaO)..... 4.7%
Magnesium (MgO) 1.8%
Boron (B)..... 0.3%

Batch Number:

Nett content: **1.0 Liter**

Registered by:



11174 Centro, Mayondon
Los Banos, Laguna

KEEP OUT OF REACH OF CHILDREN

Directions for Use:

CROP	Use Rate	Application Timing
Banana	50 - 75 ml/ 16 L tank load or 1-1.5 liters per hectare	Start application after transplanting and at 21-28 days interval.
Cabbage		Apply prior to onset of head formation and repeat at 14-21 days interval.
Potato		Apply at 14 DAT, at start of flowering and repeat every 21 days.
Rice		Apply at 14 DAT and every 21-28 days thereafter.
Pineapple		Apply during early vegetative stage at 21-28 days interval until inflorescence.

Apply an additional cycle of Alexin® just before an expected stressful event or immediately after (ie. drought, flooding, start of disease infection, etc.) to help the crop recover faster. Apply early in the morning or late afternoon when temperature is low to maximize absorption by the plants. Foliar application only.

Use this product in accordance with good agricultural practices which include the use of appropriate spray equipment set for proper coverage.

Alexin® should not be added with any sulfur or phosphite-base products.

MIXING INSTRUCTIONS:

Dilute the intended amount of Alexin® in 1 liter water container, pour solution into the spray tank and then slowly add the remaining required water volume as indicated in the above table. Mix thoroughly before application.

STORAGE AND DISPOSAL: Store in original container, tightly closed in a safe place away from children and animals. Keep in room temperature away from sunlight or any heat or hot surface.

RINSE THE CONTAINER THREE (3) TIMES AND PUNCTURE BEFORE DISPOSAL following local regulations. DO NOT REUSE EMPTY CONTAINER.

PROHIBITION AND WARRANTY:

It is a violation of FPA Rules and Regulations to use this product not in accordance with its label. Buyer assumes all risks and responsibility for safety when product is used not in accordance with directions as approved by the FPA.

Manufactured by:



South Africa

www.aeciph.com

Alexin® is a registered trademark of Nulandis® (A division of AECI Limited)