

SEE ENCLOSED LEAFLET FOR FULL DETAILS

ALEXIN®

ZEMA Reg. No.

FERTILIZER GROUP 2

A liquid organic nutrient complex containing salicylic acid.

COMPOSITION:

Calcium	(Ca)	26.4 g/kg	(33.3 g/l)
Magnesium	(Mg)	8.4 g/kg	(10.6 g/l)
Potassium	(K)	45 g/kg	(56.7 g/l)
Boron	(B)	2.375 g/kg	(3.0 g/l)

Hazard Statement

H319 Causes serious eye irritation.

Precautionary Statements

P264 Wash thoroughly after handling.

P280 Wear protective gloves/ protective clothing/ eye protection/ face protection.

P305 + P351 + P338 IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Manufacturer & Supplier:

Nulandis, a division of AECI LTD
14 Field Road, Lilianton, Boksburg, 1459,
South-Africa
Tel: +27 11 823 8000



Local Distributor:

ATS Agrochemicals Ltd.
Stand 3143, Mukwe Road, Lusaka,
Zambia
Tel: +260 1 246 724-6

ALEXIN® is a registered trademark of AECI Limited
(Nulandis®, a division of AECI Ltd)



Net Volume:

UN No. Not regulated

Batch No:

Manufacturing date:

Expiry Date:



P7043



ALEXIN®

FERTILIZER GROUP 2

A liquid organic nutrient complex containing salicylic acid.

COMPOSITION:

Calcium	(Ca)	26.4 g/kg	(33.3 g/l)
Magnesium	(Mg)	8.4 g/kg	(10.6 g/l)
Potassium	(K)	45 g/kg	(56.7 g/l)
Boron	(B)	2.375 g/kg	(3.0 g/l)

Manufacturer & Supplier:

Nulandis, a division of AECI LTD
14 Field Road, Lilianton, Boksburg, 1459,
South-Africa
Tel: +27 11 823 8000

ALEXIN® is a registered trademark of AECI Limited (Nulandis®, a division of AECI Ltd)

DIRECTIONS FOR USE: USE ONLY AS INDICATED

- Alexin should not be added with any sulphur or phosphite based products

APPLICATION TABLE:

CROP	APPLICATION RATE	APPLICATION METHOD
APPLES, PEARS	200-300 ml/100 l water to a maximum of 3 l/ha	Apply 3-4 applications from 50% flowering and repeat with 21 day intervals. For use on Pears: Do not combine Alexin with other foliar feeds or adjuvants.
CITRUS	200-300 ml/100 l water to a maximum of 3 l/ha	Apply 3 – 4 applications from 100 % petal drop and repeat with 28 day intervals.
CUCURBITS (PUMPKINS, WATERMELONS, MELONS AND CUCUMBERS)	500 ml /100 l water to a maximum of 4 l/ha	Apply at flowering and repeat with 21 day intervals.
LETTUCE, CAULIFLOWER, CABBAGE, CHINESE CABBAGE	500 ml/100 l water to a maximum of 3 l/ha	Begin applications prior to onset of head formation and repeat at 14 to 21 day intervals.
PEACHES, NECTARINES, PLUMS, APRICOTS, ALMONDS, OLIVES	200-300 ml/100 l water to a maximum of 3 l/ha	Apply 3-4 applications from 75% petal drop and repeat with 21 to 28 day intervals. If weather conditions favour bacterial infections, additional applications can be considered.
POTATOES	500 ml/100 l water to a maximum of 4 l/ha	Apply at tuber initiation and repeat every 21 to 28 days.
TOMATOES, PEPPERS, BEANS, PEAS, ONIONS	500 ml/100 l water to a maximum of 4 l/ha	Begin applications prior to fruit set and repeat at 14 to 21 day intervals.
TOBACCO	500 ml/100 l water to a maximum of 4 l/ha	Apply 2 – 3 applications bi-weekly, starting at 30 cm height.

IMPORTANT: BEFORE APPLYING ALEXIN, CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURAL ADVISER ON THE USE THEREOF.