

BZM

(POWDER)

Reg.No. B2892 Act No. 36 of 1947

FERTILIZER GROUP 2

A chelated foliar feeding mixture for correction of deficiency symptoms of crops as listed.

ACTIVE INGREDIENTS:

Boron (B) 94 g/kg
Zinc* (Zn) 43 g/kg
Manganese* (Mn) 18.5 g/kg

*EDTA chelated microelements

REGISTERED AND MANUFACTURED BY:

Gouws and Scheepers Pty (Ltd)
Reg No./ Nr.1954/003213/07
Trading as PLAASKEM
PO Box 14418
WITFIELD
1467

BZM Powder is an EDTA chelated foliar feed mixture for the correction of deficiency symptoms and for optimal growth of field crops such as tomatoes, lettuce and peppers etc.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	APPLICATION RATE	REMARKS
FIELD CROPS: Tomatoes, Lettuce and Peppers	175 g/100 l water to a maximum of 1,75 kg/ha	Apply as soon as deficiency symptoms are noticed with 7 – 10 day intervals, until no deficiency symptoms on new growth is observed
VINES	1.75 kg / ha (1 000 lt water)	Apply as soon as deficiency symptoms are noticed. Repeat with weekly intervals until no deficiency symptoms on new growth is observed. Do not apply during flowering.

BZM

(POWDER)

Reg. Nr. B2892 Wet Nr. 36 van 1947

MISSTOF GROEP 2

‘n Cheleerde blaarvoedingsmengsel vir die opheffing van tekortsimptome van gewasse soos gelys.

AKTIEWE BESTANDDELE:

Boor (B) 94 g/kg
Sink* (Zn) 43 g/kg
Mangaan* (Mn) 18.5 g/kg

*EDTA gecheleerde mikro-elemente

GEREGISTREER EN VERVAARDIG DEUR:

Gouws and Scheepers (EDMS (Bpk)
Reg No./ Nr.1954/003213/07
Handeldrywend as PLAASKEM
Posbus 14418
WITFIELD
1467

BZM Powder is ‘n EDTA gecheleerde blaarvoedingsmengsel vir die opheffing van tekortsimptome en vir optimale groei van veldgewasse, soos tamaties, slaai en groenrissies.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aanbeveel

GEWAS	DOSIS	OPMERKINGS
VELDGEWASSE: Tamaties, Slaai, Groenrissies	175 g/100 ℓ water tot ‘n maksimum van 1,75 kg/ha	Dien toe sodra tekortsimptome opgemerk word, met intervalle van 7- 10 dae, totdat geen tekortsimptome op die nuwe groei waargeneem word nie.
WINGERD	1.75 kg / ha (1 000 lt water)	Dien toe sodra tekort simptome opgemerk word. Herhaal met weeklikse intervalle, totdat geen tekort simptome op nuwe groei waargeneem word nie. Moenie gedurende die blom-kompleks spuit nie.