

CAL-NIT

CALCIUM NITRATE W/S

Reg. No. K2415, Act No. 36 of 1947
Namibian Reg. No. N-F 1463
Botswana Reg. No. W130917

FERTILIZER GROUP 1

CAL-NIT is recommended for use to alleviate calcium deficiencies.

COMPOSITION:

Calcium	(Ca)	190 g/kg
Nitrogen	(N)	155 g/kg

DIRECTIONS FOR USE:

Before applying **CAL-NIT** consult a qualified Agronomist or a Crop Adviser on the necessity of using it.

Batch No:

UN No: Not regulated

CONTENTS: **1 kg; 5 kg; 10 kg; 20 kg; 25 kg; 50 kg ; 1 ton**

Registration Holder:

Nulandis® (A division of AECI Limited)

Reg. No. 1924/002590/06

P.O. Box 14418

Witfield, 1467

Tel : (011) 823 8000

14 Field Road (Cnr Bird Street, Lilianton, 1459) Tel: (011) 823-8000

www.aeciworld.com/ www.aeciph.com



WARNINGS:

CAL-NIT may cause leaf damage if applied under hot humid conditions. Do not mix **CAL-NIT** with any other tank mixtures, apply on its own only.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED**COMPATIBILITY:**

A physical compatibility test must always be performed before such a tank mixture is sprayed.

For more information on compatibility or in the event of uncertainty, contact your local AECI Plant Health Crop Adviser.

It is recommended to do a compatibility test on a small area and observe the treated area for 3 days before doing commercial applications.

APPLICATION TABLE:

CROP	APPLICATION RATE / 100 ℓ WATER	RECOMMENDATIONS
APPLES	650 g (20 kg/ha)	Apply 6 to 8 sprays at weekly intervals from December onwards.
COTTON	500 g (2 kg/ha) 1 kg (4 kg/ha)	Seedlings: Apply at germination and repeat 14 days later. Mature plants: Apply 4 applications with 14 day intervals from flower bud formation.
CUCURBITS	75 g (300 g/ha)	Apply 3 applications with 14 day intervals from flowering.
CITRUS	400 g (10-20 kg/ha)	Apply 3 to 5 sprays with 14 day intervals from end of November.
GROUNDNUTS	1 kg (5 kg/ha)	Apply prior to flowering.
POTATOES	500 g (2,5 kg/ha)	Apply 4 sprays with 14 day intervals from flowering.
TOMATOES	350 g (2,5 kg/ha)	Apply throughout the flowering and setting stage with 14 day intervals.

CAL-NIT

KALSIUM NITRAAT W/S

Reg. Nr. K2415, Wet Nr. 36 van 1947
Namibiese Reg. Nr. N-F 1463
Botswana Reg. Nr. W130917

MISSTOF GROEP 1

CAL-NIT word gebruik om 'n tekort aan kalsium te voorkom.

SAMESTELLING:

Kalsium	(Ca)	190 g/kg
Stikstof	(N)	155 g/kg

GEBRUIKSAANWYSINGS:

Vóór toediening van **CAL-NIT** raadpleeg 'n gekwalifiseerde Agronoom of Gewasadviseur oor die noodsaaklikheid vir die gebruik daarvan.

Lotnommer:

VN Nr: Nie gereguleer nie

INHOUD: 1 kg; 5 kg; 10 kg; 20 kg; 25 kg; 50 kg ; 1 ton

Registrasiehouer:

Nulandis® ('n Afdeling van AECI Beperk)

Reg. Nr. 1924/002590/06

Posbus 14418

Witfield, 1467

Tel : (011) 823 8000

Fieldweg 14 (h/v Birdstraat, Lilianton, 1459)

Tel: (011) 823-8000

www.aeciworld.com/www.aeciph.com

WAARSKUWINGS:

CAL-NIT mag blaarskade onder warm en vogtige weersomstandighede veroorsaak.
CAL-NIT moenie met ander tenkmengsels gemeng word nie, maar slegs as 'n enkelprodukt toegedien word.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

VERENIGBAARHEID:

'n Fisiese mengbaarheidstoets eers uitgevoer word, voordat enige tenkmengsel gespuit word. Vir verdere besonderhede rakende die mengbaarheid óf in die geval van onsekerheid, kontak u plaaslike AECI Plant Health Gewasadviseur.

Dit word egter aanbeveel dat 'n toets op 'n klein area gedoen en gemonitor word vir 3 dae, voordat enige kommersiële toedienings gedoen word.

TOEDIENINGSTABEL:

GEWAS	TOEDIENINGSAANBEVELING /100 l WATER	OPMERKINGS
APPELS	650 g (20 kg/ha)	Dien 6 tot 8 bespuitings weekliks toe vanaf Desember.
KATOEN	500 g (2 kg/ha) 1 kg (4 kg/ha)	Saailinge: Dien toe met opkom en herhaal 2 weke later. Volwasse plante: Dien 4 bespuitings vanaf bolvorming toe met 14-dae tussenpose.
PAMPOEN-GEWASSE	75 g (300 g/ha)	Dien 3 bespuitings vanaf blom toe met 14-dae tussenpose.
SITRUS	400 g (10-20 kg/ha)	Dien 3 tot 5 bespuitings vanaf die einde November toe met 14-dae tussenpose.
GRONDBONE	1 kg (5 kg/ha)	Dien vóór blom toe.
AARTAPPELS	500 g (2,5 kg/ha)	Dien 4 bespuitings vanaf blomstadium toe met 14-dae tussenpose.
TAMATIES	350 g (2,5 kg/ha)	Dien vanaf blom en vrugvormingstadium toe met 14-dae tussenpose.