

CAL-NIT

CALCIUM NITRATE WS

Reg. No. K2415 Act 36 of 1947
Botswana Reg. No./ W130917, Act No 18 of 1999
Namibian Reg. No. N-F 1463

CAL-NIT is recommended for use to alleviate calcium deficiencies.

ACTIVE INGREDIENTS:

Calcium	(Ca)	190 g/kg
Nitrogen	(N)	155 g/kg

Fertilizer Group 1

Registered and Manufactured by

Gouws and Scheepers (Pty) Ltd
Reg. No. 1954/003213/07
Trading as Plaaskem
P.O. Box 14418
Witfield
1467

PRECAUTIONS:

CAL-NIT may cause leaf damage if applied under hot humid conditions. Do not mix CAL-NIT with any other tank mixes - apply on its own.

DIRECTIONS

FOR USE:

Use only as directed.

CROP	RATE/100P	APPLICATION METHODS
APPLES	650 g (20 kg/ha)	Apply 6-8 sprays at weekly intervals from December.
COTTON	500 g (2 kg/ha) 1k g (4 kg/ha)	Seedling: Apply at germination and repeat 14 days later. Mature plants: Apply at boll formulations and repeat at 14 day intervals from flowering.
CUCURBITS	75 g (300 g/ha)	Apply 3 sprays at 14 day intervals from flowering.
CITRUS	400 g (10-20 kg/ha)	Apply 3-5 sprays at 14 day intervals from end of november.
GROUNDNUTS	1 kg (5 kg/ha)	Apply prior to flowering.
POTATOES	500 g (2,5 kg/ha)	Apply 4 sprays at 14 day intervals from flowering.
TOMATOES	350 g (2,5 kg/ha)	Apply throughout the flowering and setting stage at 14 day intervals.

CAL-NIT

KALSIUM NITRAAT WO

Reg. Nr. K2415 Wet Nr 36 van 1947
Botswana Reg. Nr. W130917, Wet Nr.18 van 1999
Namibiese Reg. Nr. N-F 1463

CAL-NIT word gebruik om 'n gebrek aan kalsium te voorkom.

AKTIEWE BESTANDDELE:

Kalsium	(Ca)	190 g/kg
Stikstof	(N)	155 g/kg

Misstof Groep 1

GEREGISTREER EN VERVAARDIG DEUR:

Gouws en Scheepers (EDMS) Bpk
Reg. Nr. 1954/003213/07
Handeldrywend as Plaaskem
Posbus 14418
Witfield
1467

VOORSORGMAATREËLS:

CAL-NIT mag blaarskade onder warm en vogtige weeromstandighede veroorsaak.
CAL-NIT moet nie met ander oplossings gemeng word nie - dien alleenlik toe.

GEBRUIKSAANWYSINGS:

Gebruik alleenlik soos aangedui.

GEWAS	DOSIS/100P	OPMERKINGS
APPELS	650 g (20 kg/ha)	Dien 6-8 weeklikse bespuitings toe vanaf Desember.
KATOEN	500 g (2 kg/ha) 1 kg (4 kg/ha)	Saailinge: Dien toe met opkom en herhaal 2 weke later. Groot plante: Dien toe vanaf bolvorming en herhaal elke 2 weke vir 4 bespuitings.
PAMPOEN GEWASSE	75 g (300 g/ha)	Dien 3 twee-weeklikse bespuitings toe vanaf blom.
SITRUS	400 g (10-20 kg/ha)	Dien 3-5 twee weeklikse bespuitings toe vanaf einde November.
GRONDBONE	1 kg (5 kg/ha)	Dien voor blom toe.
AARTAPPELS	500 g (2,5 kg/ha)	Dien 4 twee weeklikse bespuitings toe vanaf blomstadium.
TAMATIES	350 g (2,5 kg/ha)	Dien elke twee weekliks toe vanaf blom en vrugvorming stadium.

