

Calcimax

LIQUID CALCIUM COMPLEX

A liquid calcium complex for use as a foliar nutrient on various crops to prevent or correct calcium deficiencies and related physiological disorders.

Inorganic Foliar Fertiliser

Composition:

Calcium (Ca)..... 78.5 g/kg (100 g/l)

(Carbohydrate chelate complex)

Boron(B)..... 5.08 g/kg (6.5 g/l)

pH: 5 - 8

STORAGE INSTRUCTIONS:

Store product in a segregated and approved area. Keep container in a cool, well-ventilated area. Keep container tightly closed and sealed until ready for use.

DIRECTIONS FOR SAFE DISPOSAL:

Invert empty container over the spray tank and allow draining for at least 30 sec after the flow has slowed down to a drip. Thereafter rinse empty container 3 times with a volume of water equal to at least 10% of that of the container and add the rinses to the contents of the spray tank before disposing of the container. Destroy empty container and do not use for any other purpose.

SUPPLIER:

NULANDIS, a division of AECI Limited
14 Field Road
Lilienton, Boksburg, 1459, South-Africa
Tel: +27 11 823 8000
www.nulandis.com



DISTRIBUTOR:

LACHLAN
Sunflag Industrial Park, Mombasa road
P.O Box 49470, 00100 Nairobi Kenya
Telephone +254 20 207 3912/3/4
www.lachlanafrica.com



Country of origin: South-Africa

Net Volume:

Batch No.:

Manufacturing date:

Expiry date:

UN No.: Not Regulated



CALCIMAX

LIQUID CALCIUM COMPLEX

A liquid calcium complex for use as a foliar nutrient on various crops to prevent or correct calcium deficiencies and related physiological disorders.

Composition:

Calcium (Ca)..... 78.5 g/kg (1 00 g/l)
(Carbohydrate chelate complex)
Boron(B)..... 5.08 g/kg (6.5 g/l)

SUPPLIER:

NULANDIS , a division of AECL Limited
14 Field Road
Lilienton, Boksburg, 1459, South-Africa
Tel: +27 11 823 8000
www.nulandis.com



DISTRIBUTOR:

LACHLAN
Sunflag Industrial Park, Mombasa road
P.O Box 49470, 00100 Nairobi Kenya
Telephone +254 20 207 3912/3/4
www.lachlanafrica.com



DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CALCIMAX is compatible with most commonly used foliar feeds. Do not use with products which contain high phosphorous, phosphite and sulphur levels including Lime sulphur or Bordeaux mixture. However, it is recommended to do a compatibility test on a small area and observe the treated area for 3 days before doing commercial applications. Where overhead systems are in use, avoid scheduling irrigation directly after application. In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation. Use the lower concentration for maintenance and the higher concentration for sufficient uptake.

SHAKE WELL BEFORE USE

APPLICATION TABLE:

CROP	DOSAGE/ha OR AS INDICATED	APPLICATION METHOD
CITRUS	300 - 500 ml/100 l water to a maximum of 15 l/ha	Apply as a medium cover spray in 1 000 - 3 000 l water/ha. Apply at fruit set.
POTATOES (Hollow heart, Brown spot) (improved shelf life and heat tolerance)	500 ml/100 l water to a maximum of 4 l/ha	Apply at tuber initiation followed by a further 3 sprays with 10 - 14 day intervals. Apply in no less than 500 l water/ha.
TOMATOES, PEPPERS (Blossom-end rot)	500 ml/100 l water to a maximum of 4 l/ha	Apply from first fruit set and repeat at 7 to 10 day intervals. Apply in no less than 500 l water/ha.
CUCURBITS (Watermelons, Cucumbers, etc.) (Blossom-end rot)	500 ml/100 l water to a maximum of 3 l/ha	Apply from first fruit set and repeat at 14 day intervals. Apply in no less than 500 l water/ha.
BRASSICAS LETTUCE	500 ml/100 l water to a maximum of 3 l/ha	Apply 2 - 3 sprays in a regular programme with 7 to 10 day intervals from heading. Apply in no less than 300 l water/ha. It is always recommended to use a sticer to improve efficacy.
CELERY (Black heart)	500 ml/100 l water to a maximum of 3 l/ha	Apply in a regular programme with 7 day intervals as a directed spray after establishment of transplanted seedlings. Apply in no less than 300 l water/ha.
SUBTROPICAL FRUIT AND NUTS: Mangoes, Avocados, Macadamias, Pecan nuts, Bananas, Litchi (Fruit colouring and internal fruit quality)	500 ml/100 l water to a maximum of 15 l/ha	Apply as a medium cover spray in 1 000 - 3 000 l water/ha during flowering and apply a follow-up spray, two weeks later.
ORNAMENTALS	500 ml / 100 l water to a maximum of 5 l/ha	Apply 2 - 3 applications with weekly intervals until symptoms disappear. Ensure thorough wetting of plants.

NOTE: Spray volumes will be determined by the actual tree or plant size. The dosage is dependent on the magnitude of the deficiency.

IMPORTANT: Before applying CALCIMAX consult a qualified Agronomist or an Agricultural Adviser on the necessity of using it.