

Cu-MAX

LIQUID COPPER COMPLEX

A liquid copper complex for use as a foliar spray on various crops to prevent or correct a copper deficiency.

COMPOSITION:

Copper (Cu)..... 4.1 %w/w (5.0 %w/v)

pH: 5-8

DIRECTION FOR SAFE DISPOSAL:

Invert empty container over the spray tank and allow draining for at least 30 sec after the flow has slowed down to a drip. Thereafter rinse empty container 3 times with a volume of water equal to at least 10% of that of the container and add the rinses to the contents of the spray tank before disposing of the container. Destroy empty container and do not use for any other purpose.

STORAGE INSTRUCTIONS:

Store product in a segregated and approved area. Keep container in a cool, well-ventilated area. Keep container tightly closed and sealed until ready for use.

UN No: Not Regulated



Inorganic Foliar Fertiliser

SUPPLIER:

AECI Plant Health
A division of AECI Ltd
14 Field Road
Lilienton, Boksburg 1459
South-Africa
T: +27 11 823 8000
www.aeciworld.com

DISTRIBUTOR:

LACHLAN
Sunflag Industrial Park,
Mombasa Road
P.O. Box 49470, 00100
Nairobi, Kenya
T: +254 20 207 3912/3/4
www.lachlanafrica.com



Country of origin: South-Africa

Batch No:

Manufacturing date:

Expiry date:

Net Volume:

Cu-MAX

A liquid organic copper complex for use as a foliar spray on various crops to prevent or correct a copper deficiency.

COMPOSITION:

Copper (Cu)..... 4.1 %w/w (5.0 %w/v)

MANUFACTURER & SUPPLIER:

AECI Plant Health

A division of AECI Ltd
14 Field Road
Lilianton, Boksburg 1459
South-Africa
T: +27 11 823 8000
www.aeciworld.com



DISTRIBUTOR:

LACHLAN

Sunflag Industrial Park,
Mombasa Road
P.O. Box 49470, 00100
Nairobi, Kenya
T: +254 20 207 3912/3/4
www.lachlanafrica.com



DIRECTIONS FOR USE: Use only as directed

- Cu-MAX is compatible with most commonly used pesticides and foliar feeds, except foliar feeds with high phosphorus content.
- Do not mix CU-MAX with highly alkaline materials such as Lime Sulphur or Bordeaux Mixture.
- Where overhead systems are in use, avoid scheduling irrigation directly after application. In orchards equipped with micro irrigation systems spray deposits should not be affected by irrigation.
- Use the lower concentration for maintenance, and the higher concentration for sufficient uptake.

CROP	DOSAGE RATE/ha OR AS INDICATED	APPLICATION DETAILS
FRUIT: Pomefruit, Stonefruit, Deciduous fruit, Subtropical fruit, Citrus	40 – 50 ml/100 l water to a maximum of 500 ml/ha	Apply in a minimum of 1 000- 5 000 l water/ha. Citrus: Apply before flowering in September, and repeat 3-4 weeks later if necessary. Other fruit: Apply with bud swell and again at 75% petal fall.
Wheat, Barley, Oats, Maize, Onions, Sugarbeet	150 – 200 ml/100 l water to a maximum of 500 ml/ha	Apply 1 – 2 applications. For best results, apply early in the season, when there is sufficient foliage to absorb the spray. Apply in a minimum of 200 l water/ha. Onions: Apply from golf ball size bulbs, and repeat 14 days later. Apply in a minimum of 500 l water/ha.
Pastures	500 ml - 1 l/ha	Apply early in the season before spring growth starts. Repeat between cuts of silage or hay. Note: Keep all livestock out of sprayed pasture for 2 weeks.
Conifers	1 l/ha	Apply to second generation trees by air, ULV or conventional sprayers.

NOTE: SPRAY VOLUMES AND DOSAGE APPLIED WILL BE DETERMINED BY THE ACTUAL TREE OR PLANT SIZE.

IMPORTANT: BEFORE APPLYING Cu-MAX CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURALIST ON THE NECESSITY OF USING IT.