



good chemistry

Cu-Max



ZEMA PRODUCT NO:

A liquid organic copper complex for use as a foliar spray on various crops to prevent or correct a copper deficiency.

COMPOSITION:

Copper (Cu).....50 g/l

Batch No:

Manufacturing date:

Expiry date:

NET VOLUME:

Manufacturer & Supplier:

AECI Plant Health

(a division of AECI Ltd.)

14 Field Road, Lilianton,

Boksburg, 1459, South-Africa

T: +27 11 823 8000

UN No: Not regulated

Read the label carefully before opening.

GROUP 2 FERTILIZER

HAZARD STATEMENTS:

Causes skin irritation.

Causes serious eye irritation.

Toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS:

Wash thoroughly after handling.

Wear protective gloves/ protective clothing/ eye protection/ face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Local Distributor:

ATS Agrochemicals Ltd.

Plot 397A, Makeni

Kafue Road, Lusaka

Zambia

T: +260 1 246 724-6



WARNING



DIRECTIONS FOR USE: Use only as directed

- Cu-MAX is compatible with most commonly used pesticides and foliar feeds, except foliar feeds with high phosphorus content.
- Do not mix CU-MAX with highly alkaline materials such as Lime Sulphur or Bordeaux Mixture.
- Where overhead systems are in use, avoid scheduling irrigation directly after application. In orchards equipped with micro irrigation systems spray deposits should not be affected by irrigation.
- Use the lower concentration for maintenance, and the higher concentration for sufficient uptake.

CROP	DOSAGE RATE/ha OR AS INDICATED	APPLICATION DETAILS
FRUIT: Pomefruit, Stonefruit, Deciduous fruit, Subtropical fruit, Citrus	40 – 50 ml/100 ℓ water to a maximum of 500 ml/ha	Apply in a minimum of 1 000- 5 000 ℓ water/ha. Citrus: Apply before flowering in September, and repeat 3-4 weeks later if necessary. Other fruit: Apply with bud swell and again at 75% petal fall.
Wheat, Barley, Oats, Maize, Onions, Sugarbeet	150 – 200 ml/100 ℓ water to a maximum of 500 ml/ha	Apply 1 – 2 applications. For best results, apply early in the sea- son, when there is sufficient foliage to absorb the spray. Apply in a minimum of 200 ℓ water/ha. Onions: Apply from golf ball size bulbs, and repeat 14 days later. Apply in a minimum of 500 ℓ water/ha.
Pastures	500 ml - 1 ℓ/ha	Apply early in the season before spring growth starts. Repeat between cuts of silage or hay. Note: Keep all livestock out of sprayed pasture for 2 weeks.
Conifers	1 ℓ/ha	Apply to second generation trees by air, ULV or conventional sprayers.

NOTE: SPRAY VOLUMES AND DOSAGE APPLIED WILL BE DETERMINED BY THE ACTUAL TREE OR PLANT SIZE.

IMPORTANT: BEFORE APPLYING Cu-MAX CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURALIST ON THE NECESSITY OF USING IT.