

DOUBLE MAX

Reg. No. B2916 Act No. 36 of 1947
Namibian Reg. No. N-F 0673

Reg. Nr. B2916 Wet Nr. 36 van 1974
Namibiese Reg. Nr. N-F 0673

FERTILIZER GROUP 2

A liquid organic calcium and-magnesium complex for correction of deficiency symptoms of crops as listed.

Active ingredients / Aktiewe bestanddele

Calcium	(Ca)	80 g / ℓ
Magnesium	(Mg)	15 g / ℓ
Boron	(B)	6.3 g / ℓ

MISSTOF GROEP 2

'n Vloeibare organiese kalsium- en magnesiumkompleks vir die opheffing van tekortsimptome van gewasse soos gelys.

Registered and Manufactured by:/Geregistreer en Vervaardig deur:

GOUWS AND SCHEEPERS (PTY) LTD / (EDMS) BPK.


REG. NO. / NR. 1954/003213/07

Trading / Handeldrywend as PLAASKEM

P.O. BOX / POSBUS 14418

WITFIELD

2041



DOUBLE MAX is a liquid organic calcium and magnesium complex for use as a foliar nutrient on various crops to prevent or correct calcium and magnesium deficiencies.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

- ❖ DOUBLE MAX is compatible with most commonly used pesticides and foliar feeds, except foliar feeds with a high phosphorous content.
- ❖ Do not Mix DOUBLE MAX with highly alkaline materials such as Lime Sulphur or Bordeaux Mixture.
- ❖ Where overhead systems are in use, avoid scheduling irrigation directly after application. In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation.
- ❖ Use the lower concentration for maintenance, and the higher concentration for sufficient uptake.
- ❖ Do not use in combination with **BLADBUFF 5**

CROP	APPLICATION RATE	REMARKS
GRAPES	500ml/100lt water to a maximum of 5lt/ha	Apply first spray when fruit has reached pea size. Repeat three times with 14 day intervals.
CITRUS	500ml/100lt water to a maximum of 10lt/ha	Apply as a medium cover spray in 1000 – 3000lt water/ha. Apply first application 100% petal drop. Repeat 1-2 times if necessary.
POME-FRUIT: Apples, Pears	500ml/100lt water to a maximum of 5lt/ha	Apply as a high volume spray in 1000 – 3000lt water/ha, depending on the tree size. Start application mid- November and continue till January / February at 7-10 day intervals.
STONE-FRUIT Apricots, Plums, Peaches	500ml/100lt water to a maximum of 8lt/ha	Apply as a medium cover spray from first week in November in 1000-3000lt water/ha. Apply a total of 6 applications with 14 day intervals.
SUB-TROPICAL FRUIT: Mangoes, Guavas, Litchis, Macadamias	500ml/100lt water to a maximum of 15lt/ha	Apply during flowering as a medium cover spray in 1000-3000lt water/ha. Apply a follow-up spray 14 days later.
POTATOES	500ml/100lt water to a maximum of 4lt/ha. Aerial application: 4lt/ha	Apply first spray with tuber initiation. Repeat every 14 to 21 days.
VEGETABLES AND FIELD CROPS: Cole crops, cucumbers, celery, peas, peppers, onions, lettuce, tomatoes, maize	500ml/100lt water to a maximum of 4lt/ha	Apply from first fruit set or heading and repeat at 7-10 day intervals. Apply in 300-500lt water/ha.
GROUNDNUTS	500ml/100lt water to a maximum of 3lt/ha	Apply in 300-500lt water/ha. Aerial application: 30-40lt water/ha. Apply early in the season when plants have sufficient leaves to absorb the spray.
PAPRICA	500ml/100lt water to a maximum of 3lt/ha	Apply when deficiency symptoms appear. Repeat if necessary. Apply in 300-500lt water/ha. Aerial application: 30-40lt water/ha.

NOTE:

SPRAY VOLUMES WILL BE DETERMINED BY THE ACTUAL TREE OR PLANT SIZE. THE DOSAGE IS DEPENDANT ON THE MAGNITUDE OF THE DEFICIENCY.

IMPORTANT: BEFORE APPLYING DOUBLE MAX CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURALIST ON THE NECESSITY OF USING IT.

DOUBLE MAX is 'n vloeibare organiese kalsium en magnesiumkompleks vir gebruik as 'n blaarvoeding op verskillende gewasse om kalsium en magnesium tekorte te voorkom of reg te stel.

GEBRUIKAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI:

DOUBLE MAX is verenigbaar met die meeste algemeen gebruikte plaagdoders en misstowwe, behalwe blaarvoedingsmiddels met 'n hoë fosfaatinhoud.

Vermyn mengsels van DOUBLE MAX en sterk alkaliese materiale soos Kalk Swawel en Bordeaux mengsel.

Waar oorhoofse besproeiing gebruik word, vermyn besproeiing van behandelde gewasse direk na toediening. In boorde waar 'n mikrobefproeiingstelsel gebruik word behoort spuitneerslae nie geaffekteer te word nie.

Gebruik die laer dosis vir instandhouding, en die hoër dosis vir die voldoende opname.

Moet nie saam met **BLADBUFF 5** gebruik nie.

GEWAS	DOSIS	OPMERKINGS
TAFELDRIUWE	500ml/100lt water tot 'n maksimum van 5lt/ha	Dien eerste bespuiting toe wanneer vrugte ertjekorrel grootte bereik het. Herhaal elke 14 dae totdat 3 bespuitings toegedien is.
SITRUS	500ml/100lt water tot 'n maksimum van 10lt/ha	Dien as 'n medium dekbespuiting to in 1000-3000lt water/ha. Dien eerste bespuiting toe met 100% blomblaarval. Herhaal 1-2 keer indien nodig.
KERNVRUGTE: Appels, Pere	500ml/100lt water tot 'n maksimum van 5lt/ha	Dien as 'n hoë volume bespuiting in 1000-3000lt water/ha toe, afhangende van die boomgrootte. Begin toediening middel November en herhaal tot Januarie / Februarie met intervale van 7 – 10 dae.
STEENVRUGTE Appelkose, Pruime, Perskes	500ml/100lt water tot 'n maksimum van 8lt/ha	Dien as 'n medium dekbespuiting toe, vanaf die eerste week in November, in 1000-3000lt water/ha. Dien 6 bespuitings in totaal toe met intervale van 14 dae.
SUB-TROPIESE VRUGTE Mango's, Guavas, Lietsjies, Makadamias	500ml/100lt water tot 'n maksimum van 15lt/ha	Dien gedurende blom toe in 1000-3000lt water/ha. Dien 'n opvolg bespuiting 14 dae later toe.
AARTAPPELS	500ml/100lt water tot 'n maksimum van 4lt/ha Lugtoediening: 4lt/ha	Dien eerste bespuiting to by knolinisasie. Herhaal elke 14 tot 21 dae.
GROENTE EN VELDGEWASSE: Kool- en pampoen gewasse, seldery, ertjies, rissies, uie, slaai, tamaties, mielies	500ml/100lt water tot 'n maksimum van 4lt/ha	Dien toe vanaf eerste vrugset of kopvorming en herhaal met intervale van 7-10 dae. Dien toe in 300-500lt water/ha.

GEWAS	DOSIS	OPMERKINGS
GRONDBONE	500ml/100lt water tot 'n maksimum van 3lt/ha	Dien toe in 300-500lt water/ha. Lugbespuiting: 30-40lt water/ha. Dien vroeg in die seisoen toe as die plant volgeodnde blare het om die bespuiting te absorbeer.
PAPRIKA	500ml/100lt water tot 'n maksimum van 3lt/ha	Dien toe sodra tekortsimptome verskryn. Herhaal indien nodig. Dien toe in 300-500lt water/ha. Lugtoediening: 30-40lt water/ha.

LET WEL: PUITVOLUMES WORD BEPAAL DEUR DIE WERKLIKE BOOM OF PLANTGROOTTE. DIE DOSIS HANG AF VAN DIE GRAAD VAN DIE TEKORT.

BELANGRIK: VOORDAT DOUBLE MAX TOEGEDIEN WORD RAADPLEEG 'N GEKWALIFISEERDE AGRONOOM OF LANDBOUKUNDIGE OOR DIE NOODSAAKLIKHEID DAARVAN.