

DOUBLE MAX

Reg. No. B2916 Act No. 36 of 1947
Namibia Reg. No. N-F 0673

Reg. Nr. B2916 Wet Nr. 36 van 1974
Namibia Reg. Nr. N-F 0673

FERTILIZER GROUP 2

A liquid organic calcium and-magnesium complex for correction of deficiency symptoms of crops as listed.

MISSTOF GROEP 2

'n Vloeibare organiese kalsium- en magnesium kompleks vir die opheffing van tekort simptome van gewasse soos aangedui.

ACTIVE INGREDIENTS / AKTIEWE BESTANDDELE:

Calcium	(Ca)	80 g / ℓ	Kalsium
Magnesium	(Mg)	15 g / ℓ	Magnesium
Boron	(B)	6.3 g / ℓ	Boor

Batch No:

Date of Manufacture:

UN No.:



REGISTRATION HOLDER: / REGISTRASIEHOUER

Nulandis® (A division of AECL Limited)' ('n Afdeling van AECL Beperk)

Reg. No./Nr 1924/002590/06
P.O. Box / Posbus 14418
Witfield, 1467
Tel : (011) 823 8000
www.nulandis.com

DOUBLE MAX is a liquid organic calcium and magnesium complex for use as a foliar nutrient on various crops to prevent or correct calcium and magnesium deficiencies.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

DOUBLE MAX is compatible with foliar feeds, except foliar feeds with a high phosphorous content. It is recommended to do a compatibility test on a small area and observe the treated area for 3 days before doing commercial applications.

- ❖ Do not mix **DOUBLE MAX** with highly alkaline materials such as Lime Sulphur or Bordeaux Mixture.
- ❖ Where overhead systems are in use, avoid scheduling irrigation directly after application. In orchards equipped with micro irrigation systems, spray deposits should not be affected by irrigation.
- ❖ Use the lower concentration for maintenance, and the higher concentration for sufficient uptake.

❖ Do not use in combination with **BLADBUFF 7** or **BLADBUFF 12**.

APPLICATION TABLE:

CROP	APPLICATION RATE	REMARKS
GRAPES	500 ml/ 100 ℓ water to a maximum of 5 ℓ/ha	Apply first spray when fruit has reached pea size stage. Repeat three times with 14 day intervals.
CITRUS	500 ml/ 100 ℓ water to a maximum of 10 ℓ/ha	Apply as a medium cover spray in 1 000 – 3 000 ℓ water/ha. Apply first application 100% petal drop. Repeat 1-2 times if necessary.
POME FRUIT: Apples, Pears	500 ml/ 100 ℓ water to a maximum of 5 ℓ/ha	Apply as a high volume spray in 1 000 – 3 000 ℓ water/ha, depending on the tree size. Start application mid- November and continue till January / February at 7-10 day intervals.
STONE FRUIT Apricots, Plums, Peaches	500 ml/ 100 ℓ water to a maximum of 8 ℓ/ha	Apply as a medium cover spray from first week in November in 1 000 – 3 000 ℓ water/ha. Apply a total of 6 applications with 14 day intervals.
SUB-TROPICAL FRUIT: Mangoes, Guavas, Litchis, Macadamias	500 ml/ 100 ℓ water to a maximum of 15 ℓ/ha	Apply during flowering as a medium cover spray in 1 000 – 3 000 ℓ water/ha. Apply a follow-up spray 14 days later.
POTATOES	500 ml/ 100 ℓ water to a maximum of 4 ℓ/ha. Aerial application: 4 ℓ/ha	Apply first spray with tuber initiation. Repeat every 14 to 21 days.
VEGETABLES AND FIELD CROPS: Cruciferae, Cucumbers, Celery, Peas, Peppers, Onions, Lettuce, Tomatoes, Maize	500 ml/ 100 ℓ water to a maximum of 4 ℓ/ha	Apply from first fruit set or heading and repeat at 7-10 day intervals. Apply in 300 – 500 ℓ water/ha.
GROUNDNUTS	500 ml/ 100 ℓ water to a maximum of 3 ℓ/ha	Apply in 300 – 500 ℓ water/ha. Aerial application: 30 – 40 ℓ water/ha. Apply early in the season when plants have sufficient leaves to absorb the spray.
PAPRIKA	500 ml/ 100 ℓ water to a maximum of 3 ℓ/ha	Apply when deficiency symptoms appear. Repeat if necessary. Apply in 300 – 500 ℓ water/ha. Aerial application: 30 – 40 ℓ water/ha.

NOTE:

SPRAY VOLUMES WILL BE DETERMINED BY THE ACTUAL TREE OR PLANT SIZE.

THE DOSAGE IS DEPENDANT ON THE MAGNITUDE OF THE DEFICIENCY.

IMPORTANT: BEFORE APPLYING DOUBLE MAX CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURALIST ON THE NECESSITY OF USING IT.

BLADBUFF® 7 and BLADBUFF 12® are the registered trademarks of AECI Limited (Nulandis® a division of AECI Ltd.)

DOUBLE MAX is 'n vloeibare organiese kalsium- en magnesiumkompleks vir gebruik as 'n blaarvoeding op verskillende gewasse om kalsium en magnesium tekorte te voorkom of reg te stel.

GEBRUIKAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

DOUBLE MAX is verenigbaar met misstowwe, behalwe blaarvoedingsmiddels met 'n hoë fosfaatinhoud. Dit word egter aanbeveel dat 'n toets op 'n klein area gedoen word en dan nagegaan word vir 3 dae, voordat enige kommersiële toedienings gedoen word.

Vermyn mengsels van **DOUBLE MAX** en sterk alkaliese materiale soos Kalk Swael en Bordeaux mengsel.

Waar oorhoofse besproeiing gebruik word, vermyn besproeiing van behandelde gewasse direk na toediening. In boorde waar 'n mikrobefproeiingstelsel gebruik word behoort spuitneerslae nie geaffekteer te word nie.

Gebruik die laer dosis vir instandhouding, en die hoër dosis vir voldoende opname.

Moenie saam met **BLADBUFF 7** of **BLADBUFF 12** gebruik nie.

TOEDIENINGSTABEL:

GEWAS	DOSIS	OPMERKINGS
TAFELDRUIWE	500 ml/ 100 ℓ water tot 'n maksimum van 5 ℓ/ha	Dien eerste bespuiting toe wanneer vrugte ertjekorrelgrootte stadium bereik het. Herhaal elke 14 dae totdat 3 bespuitings toegedien is.
SITRUS	500 ml/ 100 ℓ water tot 'n maksimum van 10 ℓ/ha	Dien as 'n medium dekbespuiting toe in 1 000 – 3 000 ℓ water/ha. Dien eerste bespuiting toe met 100% blomblaarval. Herhaal 1 tot tweekeer indien nodig.
KERNVRUGTE: Appels, Pere	500 ml/ 100 ℓ water tot 'n maksimum van 5 ℓ/ha	Dien as 'n hoë volume bespuiting in 1 000 – 3 000 ℓ water/ha toe, afhangende van die boomgrootte. Begin toediening middel November en herhaal tot Januarie / Februarie met intervale van 7 tot 10 dae.
STEENVRUGTE Appelkose, Pruime, Perskes	500 ml/ 100 ℓ water tot 'n maksimum van 8 ℓ/ha	Dien as 'n medium dekbespuiting toe vanaf die eerste week in November, in 1 000 – 3 000 ℓ water/ha. Dien 6 bespuitings in totaal toe met intervale van 14 dae.
SUBTROPIESE VRUGTE Mango's, Guavas, Lietsjies, Makadamias	500 ml/ 100 ℓ water tot 'n maksimum van 15 ℓ/ha	Dien gedurende blom toe in 1 000 tot 3 000 ℓ water/ha. Dien 'n opvolg bespuiting 14 dae later toe.
AARTAPPELS	500 ml/ 100 ℓ water tot 'n maksimum van 4 ℓ/ha <u>Lugtoediening:</u> 4 ℓ/ha	Dien eerste bespuiting toe tydens knolinisasie. Herhaal elke 14 tot 21 dae.

GROENTE EN VELDGEWASSE: Kool- en Pampoen gewasse, Seldery, Ertjies, Rissies, Uie, Slaai, Tamaties, Mielies	500 m ³ / 100 ℓ water tot 'n maksimum van 4 ℓ/ha	Dien toe vanaf eerste vrugset of kopvorming en herhaal met intervalle van 7 tot 10 dae. Dien toe in 300 – 500 ℓ water/ha.
GRONDBONE	500 m ³ / 100 ℓ water tot 'n maksimum van 3 ℓ/ha	Dien toe in 300 – 500 ℓ water/ha. Lugtoediening: 30-40 ℓ water/ha. Dien vroeg in die seisoen toe as die plant voldoende blare het om die bespuiting te absorbeer.
PAPRIKA	500 m ³ / 100 ℓ water tot 'n maksimum van 3 ℓ/ha	Dien toe sodra tekort simptome verskyn. Herhaal indien nodig. Dien toe in 300 – 500 ℓ water/ha. Lugtoediening: 30-40 ℓ water/ha.

**LET WEL: PUITVOLUMES WORD BEPAAL DEUR DIE WERKLIKE BOOM
OF PLANTGROOTTE. DIE DOSIS HANG AF VAN DIE GRAAD VAN DIE TEKORT.**

**BELANGRIK: VOORDAT DOUBLE MAX TOEGEDIEN WORD RAADPLEEG 'N
GEKWALIFISEERDE AGRONOOM OF LANDBOUKUNDIGE OOR DIE
NOODSAAKLIKHEID DAARVAN.**

BLADBUFF® 7 en BLADBUFF 12® is die geregistreerde handelsmerke van AECI Beperk
(Nulandis® 'n afdeling van AECI Beperk)