

Main Panel

Eradicate Plus 720 EC

Reg.No./Nr.L5249, Act No. / Wet Nr. 36 of/van 1947
Namibian/Namibiese Reg. No./Nr. N-AR 1319

A SELECTIVE SOIL INCORPORATED EMULSIFIABLE CONCENTRATE HERBICIDE FOR THE CONTROL OF SEDGES (UINTJIES), ANNUAL GRASSES AND CERTAIN BROADLEAF WEEDS IN CROPS AS INDICATED.

'n SELEKTIEWE GRONDINGEWERKTE, EMULGEERBARE KONSENTRAAT ONKRUIDDODER VIR DIE BEHEER VAN UINTJIES, EENJARIGE GRASSE EN SEKERE BREëBLAARONKRUIDE IN GEWASSE SOOS AANGEDUI.

HERBICIDE GROUP CODE N ONKRUIDDODERGROEPKODE

ACTIVE INGREDIENT: / AKTIEWE BESTANDDEEL :

EPTC (Thiocarbamate).....720 g/l... ... EPTC (Tiokarbamaat)
Contains Dichlormid (Safener) 66.4 g/lBevat Diklormied (Beveiliger)

Net Volume



Netto Volume



REGISTRATION HOLDER / REGISTRASIEHOUER:

Nulandis® (A division of AECI Limited) / Nulandis® ('n Afdeling van AECI Beperk)

Reg. No. / Nr. 1924/002590/06

P.O. Box / Posbus 14418

Witfield, 1467

Tel : (011) 823 8000

www.nulandis.com

IN CASE OF POISONING PLEASE PHONE / IN GEVAL VAN VERGIFTIGING SKAKEL 021 931-6129/
0824668946

Batch No:
Lotnommer:

Date of Manufacture:
Datum van Vervaardiging:

UN No. VN Nr. 3006 (Thiocarbamate Pesticide Liquid, Toxic)



HARMFUL
SKADELIK



PAMPHLET

Eradicate Plus 720 EC

Reg.No. L 5249, Act No. 36 of 1947
Namibian Reg. No. N-AR 1319

A SELECTIVE SOIL INCORPORATED EMULSIFIABLE CONCENTRATE HERBICIDE FOR THE CONTROL OF SEDGES (UINTJIES), ANNUAL GRASSES AND CERTAIN BROADLEAF WEEDS IN CROPS AS INDICATED.

HERBICIDE GROUP CODE N

ACTIVE INGREDIENT:

EPTC (Thiocarbamate).....720 g/l
Contains Dichlormid (Safener)..... 66.4 g/l

REGISTRATION HOLDER

Nulandis® (A division of AECI Limited)
Reg. No. 1924/002590/06
P.O. Box 14418
Witfield, 1467
Tel : (011) 823 8000



HARMFUL

WARNINGS:

Poisonous if swallowed.

May cause irritation to skin and eyes.

FLAMMABLE - do not store near open flames.

Store away from food and feeds.

Keep out of reach of children, uninformed person and animals.

Eradicate Plus 720 EC should only be used if soil conditions are suitable for fast germination of the crop.

Re-entry: Do not enter treated area within 1 day after treatment unless wearing protective clothing.

In case of poisoning – CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM.

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL CLIMATIC AND STORAGE CONDITIONS; QUALITY OF DILUTION WATER; COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF WEEDS AGAINST THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS:

Do not inhale spray mist or fumes. Close container after use.

Wash with soap and water after use or skin contact. In case of eye contact, rinse immediately with running water for at least 15 minutes.

Do not eat, drink or smoke whilst mixing or applying, or before washing hands and face and changing clothes.

Prevent contamination of food, feeds, drinking water and eating utensils.

Clean applicator thoroughly with **FLUSHSOL** before using it with other material and dispose of wash water where it will not contaminate crops, grazing, rivers or dams. Refer to the **FLUSHSOL** label for directions of use.

When empty, invert the container over the spray tank or mixing tank and allow it to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to at least one tenth of that of the container and add the rinsings to the contents of the spray tank. Destroy the container by perforation and flattening and do not use for any other purpose.

NOTE: ERADICATE PLUS 720 EC contains a petroleum distillate as solvent and if swallowed, do not induce vomiting but call a doctor.

SYMPTOMS OF POISONING:

Symptoms may include the following; Eye and skin irritation. Respiratory tract irritation may lead to frequent attacks of bronchial infection.

FIRST AID TREATMENT

INHALATION: If inhaled, remove patient to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Get medical attention.

INGESTION: DO NOT INDUCE vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. If large quantities of this material have been swallowed, call a physician immediately. Loosen tight clothing such as a collar, tie, belt or waistband.

SKIN CONTACT: In case of contact, immediately flush skin with plenty of water. Remove contaminated clothing and shoes. Thoroughly wash shoes and clothing before re-use. Get medical attention.

EYE CONTACT: Check for and remove any contact lenses. In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. WARM water MUST be used. Get medical attention.

NOTE TO PHYSICIAN

No specific treatment – treat symptomatically.

RESISTANCE WARNING:

For resistance management, **Eradicate Plus 720 EC** is a group code N herbicide. Any weed population may contain individuals naturally resistant to **Eradicate Plus 720 EC** and other group code N herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. The resistant weeds may not be controlled by **Eradicate Plus 720 EC** or any other group code N herbicides.

To delay herbicide resistance:

- Avoid exclusive repeated use of herbicides from the same herbicide group code. Alternate or tank mix with registered products from different herbicide group codes.
- Integrate other control methods (chemical, cultural, biological) into weed control programmes.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE : USE ONLY AS DIRECTED

USE RESTRICTIONS:

ERADICATE PLUS 720 EC must only be used when conditions are ideal for fast germination of the crop and weed.

DO NOT use on maize grown for seed before consulting the seed company.

DO NOT use as an overall application on kidney beans.

ERADICATE PLUS 720 EC is a herbicide which must be thoroughly incorporated with the upper 100 - 500 mm layer of soil before planting or it should be injected into the soil at planting with suitable equipment. Incorporation should take place immediately after application.

SOIL PREPARATION

The seed bed must be thoroughly prepared and free of unnecessary clods, plant residues and weeds. All growing weeds, especially nutsedge, must be destroyed before application.

Continuous use of **ERADICATE PLUS 720 EC** on the same lands year after year, may result in decreased herbicidal efficacy. As a precaution **ERADICATE PLUS 720 EC** should be rotated with a herbicide from a different herbicide group code on an annual basis.

METHODS OF APPLICATION:

Band Application

Only use equipment which will apply **ERADICATE PLUS 720 EC** as recommended.

Calibrate the equipment under actual field conditions before use.

Check the calibration during application from time to time.

Overall Application

For overall weed control, apply **ERADICATE PLUS 720 EC** as an overall spray to a well prepared seed bed, free of growing weeds and unnecessary clods.

Thoroughly incorporate the weed killer immediately after application or within minutes of application into the upper 100 - 500 mm soil layer. Application and incorporation should rather be carried out in one step.

Incorporation can be carried out using a tandem disc to a depth of at least 150 mm at 7 - 8 km/h followed by an inverted spike tooth harrow or other device which levels wider than the disc. For thorough mixing, disc twice (criss-cross). Prior to the second discing reset the disc to prevent cutting deeper than the first discing. Apply the recommended quantity of **ERADICATE PLUS 720 EC** in sufficient water per ha (200 ℓ - 450 ℓ/ha) to ensure an even distribution of the herbicide.

APPLICATION TABLE:

CROP	DOSAGE PER ha AND CLAY CONTENT OF SOIL		REMARKS
MAIZE, SWEETCORN, SWEET POTATOES & LUCERNE	% CLAY	DOSAGE	OVERALL APPLICATION The seed should be planted as soon as possible after application to obtain the maximum period of weed control. In the case of lucerne, wait 5 days before sowing. If it is necessary to cultivate after application and incorporation, do not work deeper than the depth to which the herbicide was incorporated. BAND APPLICATION To obtain a 450 mm band of weed control apply ERADICATE PLUS 720 EC 40 - 60 mm deep using a suitable applicator to produce four sub-surface streams 80 mm and 170 mm on either side of the sown rows or, for example, by means of a suitable sweep mounted on the planter.
	0 - 10	2,0 ℓ	
	11 - 15	2,5 ℓ	
	16 - 20	3,0 ℓ	
	21 - 25	3,5 ℓ	
	26 - 30	4,0 ℓ	
SUGARCANE	% CLAY	DOSAGE	OVERALL APPLICATION FOR PLANTCANE Apply the recommended rate of ERADICATE PLUS 720 EC as an overall application and incorporate to a depth of 200 mm as described for maize. Planting should be carried out immediately after incorporation to obtain the maximum period of weed control. Planting by hand is recommended. Plant furrows should not exceed 150 mm in depth to prevent untreated soil being brought to the surface.
	0 - 15	3 ℓ	
	16 - 30	5 ℓ	
	more than 30	7 ℓ	

CROP	DOSAGE PER ha AND CLAY CONTENT OF SOIL		REMARKS
POTATOES	% CLAY DOSAGE 0 - 15 3,5 ℓ 16 - 30 5,3 ℓ		OVERALL APPLICATION Apply the recommended dosage ERADICATE PLUS 720 EC as an overall spray and incorporate as described for maize. Hereafter make furrows and plant as normal. BAND APPLICATION Place fertilizer and tubers in position in open furrows and cover with a 50 mm layer of soil. Ensure that furrow sides are not steeper than 45° with good retention of the spray. Apply as a band over the row and close the furrows immediately to cover the ERADICATE PLUS 720 EC.
DRY BEANS <i>(Phaseolus vulgaris)</i> AND GREEN BEANS (CAPE CULTIVAR)	% CLAY DOSAGE 0 - 10 1,5 ℓ 11 - 15 2,0 ℓ 16 - 20 2,5 ℓ 21 - 30 3,0 ℓ		OVERALL APPLICATION Apply as for maize. BAND APPLICATION Apply only as sub-surface injection as for maize. Initial retardation of seedlings may occur, but plants will develop normally.
KIDNEY BEANS <i>(Phaseolus coccineus)</i>	% CLAY DOSAGE per 10 000 m row length and a band of 450 mm 0 - 10 620 ℓ 11 - 30 780 ℓ		BAND APPLICATION Only a band application is recommended. Apply only as sub-surface streams as recommended for maize and ensure that the two inner streams are 40 mm above the seed level.

CROP	DOSAGE PER ha AND CLAY CONTENT OF SOIL		REMARKS
SUNFLOWERS	% CLAY	DOSAGE	OVERALL APPLICATION Must only be used on Kortrus selections. Apply to a well prepared seedbed and incorporate immediately. To reduce initial retardation of seedlings, only plant 7 days after application.
	0 - 15	not recommended	
	16 - 20	2,0 l	
	21 - 30	3,0 l	

LIST OF WEEDS CONTROLLED BY ERADICATE PLUS 720 EC:

Botanical Name

Common Name

NUTSEDGES:

Cyperus esculentus
Cyperus rotundus

Yellow nutsedge
 Purple nutsedge

GRASSES:

Brachiara eruciformis
Chloris pycnothrix
Digitaria sanguinalis
Eleusine indica
Panicum schinzii
Setaria pallide-fusca
Setaria verticillata
Tragus racemosus
Urochloa panicoides

Sweet signal grass
 Spiderweb chloris
 Crab finger-grass
 Goose grass
 Sweet buffalo grass
 Red bristle grass
 Sticky bristle grass
 Large carrot-seed grass
 Herringbone grass

BROADLEAF WEEDS:

The control of the following weeds may vary according to conditions. They will only be controlled when the correct dosage is used in a well prepared fine seedbed followed by soil conditions favourable for germination of seed.

Amaranthus deflexus
Amaranthus hybridus
Bidens pilosa
Cleome monophylla
Chenopodium album
Datura stramonium

Perennial pigweed
 Common pigweed
 Blackjack
 Spindlepod
 White goosefoot
 Thorn apple

Flushsol® is the registered trademark of AECI Limited (Nulandis® a division of AECI Ltd.)

Pamflet

Eradicate Plus 720 EC

Reg.Nr. L5249, Wet Nr. 36 van 1947
Namibië: Reg. Nr. N-AR 1319

'n SELEKTIEWE GRONDINGEWERKTE, EMULGEERBARE KONSENTRAAT
ONKRUIDDODER VIR DIE BEHEER VAN UINTJIES, EENJARIGE GRASSE EN SEKERE
BREëBLAARONKRUIDE IN GEWASSE SOOS AANGEDUI.

ONKRUIDDODERGROEPKODE N

AKTIEWE BESTANDDEEL :

EPTC (Tiokarbamaat)..... 720 g/l
Bevat Diklormied (Beveiliger)..... 66.4 g/l

REGISTRASIEHOUER:

Nulandis® ('n Afdeling van AECI Beperk)
Reg. Nr. 1924/002590/06
Posbus 14418
Witfield, 1467
Tel : (011) 823 8000



SKADELIK

WAARSKUWINGS:

Giftig indien ingeneem.

Mag vel- en oogirritasie veroorsaak.

VLAMBAAR - moenie naby oop vlamme berg nie.

Berg weg van voedsel en voer.

Hou buite bereik van kinders, oningeligte persone en diere.

Eradicate Plus 720 EC moet slegs gebruik word indien grondtoestande geskik is vir vinnige ontkieming van die gewas.

Herbetreding: Moenie behandelde gebied betree vir 1 dag na die bespuiting nie, tensy beskermende oorklere gedra word.

In geval van vergiftiging – ONTBIED 'N GENEESHEER EN STEL HIERDIE ETIKET AAN HOM BESIKBAAR.

ALHOEWEL HIERDIE MIDDEL OMVATTEND ONDER 'N GROOT VERSKEIDENHEID TOESTANDE GETOETS IS WAARBORG DIE REGISTRASIEHOUER NIE DAT DIT ONDER ALLE TOESTANDE DOELTREFFEND SAL WEES NIE AANGESIEN DIE WERKING EN EFFEK DAARVAN BEÏNVLOED KAN WORD DEUR FAKTORE SOOS ABNORMALE KLIMAATS- EN BERGINGSTOESTANDE; KWALITEIT VAN VERDUNNINGSWATER, VERENIGBAARHEID MET ANDER STOWWE WAT NIE OP DIE ETIKET AANGEDUI IS NIE EN DIE VOORKOMS VAN WEERSTAND VAN DIE ONKRUIDE TEEN DIE BETROKKE MIDDEL SOWEL AS DIE METODE, TYD EN AKKURAATHEID VAN TOEDIENING. VERDER AANVAAR DIE REGISTRASIEHOUER NIE VERANTWOORDELIKHEID VIR SKADE AAN GEWASSE, PLANTEGROEI, DIE OMGEWING OF VIR NADELIGE EFFEK OP MENS OF DIER OF VIR 'N GEBREK AAN PRESTASIE VAN DIE BETROKKE MIDDEL AS GEVOLG VAN DIE VERSUIM DEUR DIE GEBRUIKER OM ETIKETAANWYSINGS NA TE KOM OF AS GEVOLG VAN DIE ONTSTAAN VAN TOESTANDE WAT NIE KRAGTENS DIE REGISTRASIE VOORSIEN KON WORD NIE. RAADPLEEG DIE VERSKAFFER IN DIE GEVAL VAN ENIGE ONSEKERHEID.

VOORSORGMATREËLS:

Moet nie dampe of sproeinewel inasem nie.

Was met seep en water na gebruik en toevallige kontak met vel.

Besoedelde klere moet daaglik gewas word.

Was met seep en water na gebruik of velkontak. In geval van oogkontak spoel onmiddellik vir minstens 15 minute met lopende water.

Moenie eet, rook of drink tydens vermenging of toediening of voordat hande en gesig gewas en skoon klere aangetrek is nie.

Verhoed besoedeling van voedsel, voer, drinkwater en eetgerei.

Verhoed wegdrywing van die sproeinewel na ander gewasse, weidings, riviere, damme en ander gebiede wat nie behandel word nie.

Maak toedieningsapparaat skoon met **Flushsol** voor her-gebruik en doen weg met waswater waar dit nie gewasse, weiding, riviere en damme sal besoedel nie. Verwys na die **Flushsol** etiket vir gebruiksaanwysings.

Sodra die houer leeg is keer dit om oor die spuittenk of mengbak en dreineer vir minstens 30 sekondes nadat die vloei tot 'n gedrup verminder het. Spoel die houer daarna driekeer uit met 'n volume water gelyk aan minstens 'n tiende van die houer en gooi die spoelwater by die inhoud van die spuittenk. Vernietig die houer deur gate daarin te maak en plat te druk. Moenie vir enige ander doel gebruik nie.

NOTA: **ERADICATE PLUS 720 EC** bevat 'n petroleumdistillaat as oplosmiddel en moet nie braking veroorsaak wanneer per ongeluk ingesluk word nie maar ontbied 'n geneesheer.

SIMPTOME VAN VERGIFTIGING:

Simptome mag die volgende insluit; Oog- en velirritasie. Irritasie van asemhalingsstelsel mag aanleiding gee tot gereelde brongitus infeksie.

NOODHULPBEHANDELING:

INASEMING: Indien ingeasem, verwyder pasiënt na vars lug. Indien pasiënt ophou asemhaal, pas kunsmatige asemhaling toe. Indien asemhaling bemoeilik word, gee suurstof. Verkry mediese hulp.

INGESLUK: MOENIE braking veroorsaak tensy so aanbeveel deur mediese personeel. Moet nooit iets per mond aan 'n bewustelose persoon gee nie. Indien groot hoeveelhede van die middel gedrink is skakel onmiddellik 'n geneesheer. Maak styfpasende klere, soos bv. 'n kraag, das, belt of lyfband los.

VELKONTAK: Ingeval van velkontak, spoel onmiddellik af met genoegsame water. Verwyder besoedelde klere en skoene. Besoedelde klere en skoene moet deeglik gewas word voor her-gebruik. Verkry mediese hulp.

OOGKONTAK: Gaan na en verwyder enige kontaklense. In geval van oogkontak, spoel onmiddellik met genoegsame water vir minstens 15 minute. WARM water MOET gebruik word. Verkry mediese hulp.

INLIGTING AAN GENEESHEER:

Geen spesifieke behandeling – behandel simptome.

WEERSTANDSWAARSKUWING:

Eradicate Plus 720 EC is 'n groepkode N onkruidodder. Enige populasie van 'n spesifieke onkruidspesie mag individue insluit wat 'n natuurlike weerstand teen **Eradicate Plus 720 EC** of enige ander groepkode N onkruidodders het. Indien hierdie onkruidodders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die onkruidpopulasie oorheers. Hierdie weerstandbiedende onkruid sal waarskynlik nie deur **Eradicate Plus 720 EC** of enige ander groepkode N onkruidodders beheer word nie.

Om weerstand teen onkruidodders te vertraag:

- Vermyn die eksklusiewe herhaaldelike gebruik van onkruidodders met dieselfde groepkode. Wissel af met, of gebruik tenkengsels van produkte in verskillende onkruidoddergroepkodes.
- Integreer ander beheermaatreëls (chemies, verbouing, biologies) in onkruidbeheerprogramme.

Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.

GEBRUIKSAANWYSINGS : GEBRUIK SLEGS SOOS AANBEVEEL

GEBRUIKSBEPERKINGS:

ERADICATE PLUS 720 EC moet slegs gebruik word wanneer toestande ideaal is vir vinnige ontkieming van die gewas en onkruid.

MOENIE op mielies gebruik wat vir saad gekweek word voordat die betrokke saadmaatskappy geraadpleeg is nie.

MOENIE as 'n algehele toediening op nierbone gebruik nie.

ERADICATE PLUS 720 EC is 'n onkruidodder wat net vóór plant deeglik met die boonste 100 - 150 mm grondlaag vermeng moet word voordat geplant word. Dit kan ook tydens plant met geskikte toerusting in die grond ingespuut word. Inwerking moet onmiddellik na toediening plaasvind.

VOORBEREIDING VAN GROND:

Die saadbed moet deeglik voorberei wees sonder onnodige kluite, plantreste en onkruid. Alle groeiende onkruid, veral uittjies, moet voor toediening doodgework word.

Aanhoudende gebruik van **ERADICATE PLUS 720 EC** jaar na jaar in dieselfde grond, mag aanleiding gee tot verminderde onkruidbeheer. As voorsorgmaatreël behoort **ERADICATE PLUS 720 EC** in rotasie met 'n ander gewas en groepkode onkruidodder, op 'n jaarlikse, basis gebruik te word.

METODES VAN TOEDIENING:

Strooktoediening:

Gebruik slegs 'n toediener wat **ERADICATE PLUS 720 EC** volgens die aanbevole wyse kan toedien. Kalibreer die toerusting onder werklike veldtoestande voordat dit gebruik word. Gaan die kalibrasie van tyd tot tyd in die veld na.

Algehele toediening:

Vir algehele onkruidbeheer dien **ERADICATE PLUS 720 EC** as 'n algehele bespuiting toe op 'n goed voorbereide saadbed, vry van groeiende onkruid en onnodige kluite. Werk die middel onmiddellik of enkele minute na toediening deeglik in die boonste 100 - 150 mm grondlaag. Toediening en inwerk moet liefst gelyktydig gedoen word. Inwerking kan d.m.v. 'n tandemskotteleg óf 'n wisselgangskotteleg tot op 'n diepte van ten minste 150 mm teen 7 - 8 km/h uitgevoer word, gevolg deur 'n omgekeerde tandeg of sleepyster wat breër sleep as die skotteleg. Vir deeglike vermenging bewerk twee keer (kruis en dwars) in. Voor die tweede bewerking verstel die skotteleg om te voorkom dat dit dieper as die eerste bewerking sny. Dien die aanbevole **ERADICATE PLUS 720 EC** in voldoende water per ha toe (200 - 450 ℓ/ha) om 'n eweredige verspreiding van die onkruidodder te verseker.

TOEDIENINGSTABEL:

GEWAS	DOSIS PER ha EN KLEI-INHOUD VAN GROND		OPMERKINGS
MIELIES, SUIKERMIELIES, PATATS & LUSERN	% KLEI	DOSIS	ALGEHELE TOEDIENING Die saad moet so gou as moontlik na toediening geplant word om die maksimum tydperk van onkruidbeheer te verkry. In die geval van lusern, wag 5 dae voor plant. Indien dit nodig is om na toediening en inwerking te skoffel, moet die grond nie dieper as die diepte waarop die onkruidodder ingewerk is, bewerk word nie. STROOKBEHANDELING Vir 'n onkruidvrye strook van 450 mm oor die mielierye dien die ERADICATE PLUS 720 EC 40 - 60 mm diep toe met 'n geskikte toediener - vier ondergrondse straaltpies 80 en 170 mm weerskante van die saadrye of m.b.v. 'n vlerkskaarimplement wat op die planter gemonteer is.
	0 - 10	2,0 ℓ	
	11 - 15	2,5 ℓ	
	16 - 20	3,0 ℓ	
	21 - 25	3,5 ℓ	
	26 - 30	4,0 ℓ	
SUIKERRIET	% KLEI	DOSIS	ALGEHELE TOEDIENING VIR PLANTRIET Dien die aanbevole dosis ERADICATE PLUS as 'n algehele bespuiting toe en werk in tot op 'n diepte van 200 mm soos uiteengesit vir mielies. Plant onmiddellik hierna om die maksimum tyd van onkruidbeheer te verkry. Handplanting word aanbeveel. Plantvoor diepte moenie 150 mm oorskry nie om te voorkom dat onbehandelde grond na die oppervlak gebring word.
	0 - 15	3 ℓ	
	16 - 30	5 ℓ	
	meer as 30	7 ℓ	

GEWAS	DOSIS PER ha EN KLEI-INHOUD VAN GROND		OPMERKINGS
AARTAPPELS	% KLEI	DOSIS	ALGEHELE TOEDIENING Dien die aanbevole hoeveelheid ERADICATE PLUS 720 EC toe as 'n algehele bespuiting en werk in soos verduidelik vir mielies. Trek hierna die vore en plant soos normaalweg. STROOKBEHANDELING Plaas die moere en kunsmis in posisie in die oop vore en bedek met 'n 50 mm grondlagie. Die wande van die vore moet nie steiler as 45° wees nie en moet gerieflik gespuit kan word sonder afloop. Dien as 'n strookbehandeling oor die rye toe en maak die vore onmiddellik toe om die ERADICATE PLUS 720 EC te bedek.
	0 - 15 16 - 30	3,5 ℓ 5,3 ℓ	
DROËBONE (<i>Phaseolus vulgaris</i>) EN GROENBONE (CAPE KULTIVAR)	% KLEI	DOSIS	ALGEHELE TOEDIENING Dien toe soos by mielies. STROOKBEHANDELING Dien slegs toe met ondergrondse inspuittande soos by mielies. Aanvanklike groei-vertraging mag voorkom, maar plante sal normaalweg herstel.
	0 - 10 11 - 15 16 - 20 21 - 30	1,5 ℓ 2,0 ℓ 2,5 ℓ 3,0 ℓ	
NIERBONE (<i>Phaseolus coccineus</i>)	% KLEI	DOSIS per 10 000 m rylengte en 'n strook van 450 mm	STROOKBEHANDELING Slegs 'n strookbehandeling word aanbeveel. Dien slegs toe met inspuittande soos by mielies en maak seker dat die 2 binneste straaltjies 40 mm bo die saadvlak is.
	0 - 10 11 - 30	620 ℓ 780 ℓ	

GEWAS	DOSIS PER ha EN KLEI-INHOUD VAN GROND		OPMERKINGS
SONNEBLOM	% KLEI	DOSIS	ALGEHELE TOEDIENING Moet slegs op Kortrus-seleksie gebruik word. Dien toe op 'n goed voorbereide saadbed en werk onmiddellik in. Om aanvanklike vertraging van saailinge te verminder plant 7 dae na toediening.
	0 - 15	nie aanbeveel	
	16 - 20	2,0 ℓ	
	21 - 30	3,0 ℓ	

LYS VAN ONKRUIDE WAT DEUR ERADICATE PLUS 720 EC BEHEER WORD

Botaniese Naam

Gewone Naam

UINTJIES:

Cyperus esculentus
Cyperus rotundus

Geeluintjie
Rooiuintjie

GRASSE:

Brachiara eruciformis
Chloris pycnothrix
Digitaria sanguinalis
Eleusine indica
Panicum schinzii
Setaria pallide-fusca
Setaria verticillata
Tragus racemosus
Urochloa panicoides

Litjiesinjalgras
Spinnerak chloris
Kruisvingergras
Jongosgras
Soetbuffelsgras
Rooiborselgras
Klitsborselgras
Grootwortelsaadgras
Beesgras

BREËBLAARONKRUIDE:

Die beheer van die volgende breëblaaronkruid sal wissel nagelang van omstandighede. Hulle sal slegs beheer word indien die korrekte dosis gebruik word op 'n goed voorbereide en fyn saadbed gevolg deur grondtoestande wat gunstig is vir saadontkieming.

Amaranthus deflexus
Amaranthus hybridus
Bidens pilosa
Cleome monophylla
Chenopodium album
Datura stramonium

Meerjarige misbredie
Gewone misbredie
Knapsekêrel
Rusperbossie
Withondebossie
Stinkblaar

Flushol® is die geregistreerde handelsmerk van AECl Beperk (Nulandis® 'n Afdeling van AECl Bpk.)