

# ERASE 360 SL

Reg. No./Nr. L6206, Act No./Wet Nr. 36 of/van 1947  
Botswana Reg. No./Nr. W130314, Act No./Wet Nr.18 of/van 1999  
Malawi Reg. No./Nr. MW/PCB/2008/0120  
Namibian Reg. No./Namibiese Reg. Nr. N-AR 1200



**READ THE LABEL BEFORE USE**  
**KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

A NON-SELECTIVE SYSTEMIC POST-EMERGENCE HERBICIDE FOR THE CONTROL OF PERENNIAL AND ANNUAL WEEDS IN AGRICULTURE AND INDUSTRIAL AREAS.

'n NIE-SELEKTIEWE SISTEMIESE NA-OPKOM ONKRUIDDODER VIR DIE BEHEER VAN EEN- EN MEERJARIGE ONKRUIDE IN LANDBOU EN INDUSTRIËLE GEBIEDE.

GROUP/GROEP

9

HERBICIDE/ONKRUIDDODER

**ACTIVE INGREDIENT:**

Glyphosate (Isopropylamine salt) .....480 g/ℓ.....Glifosaat(Isopropielamien sout)

Glyphosate (acid equivalent).....360 g /ℓ.....Glifosaat (suur ekwivalent)

**Net Volume**

ℓ

**Netto Volume**



**DANGER**

**GHS Hazard Statements**

Causes skin irritations.

Causes serious eye damage.

Toxic to aquatic life with long lasting effects.

**GHS Precautionary Statements**

Do not breathe dust, fume, gas, mist, vapours or spray.

Wash thoroughly after handling



**REGISTRATION HOLDER / REGISTRASIEHOUER:**

**Nulandis® (A division of AECI Limited)/ Nulandis® ('n Afdeling van AECI Beperk)**

Reg. No. / Nr. 1924/002590/06

P.O. Box / Posbus 14418

Witfield, 1467

Tel : (011) 823 8000

www.aeciph.com

IN CASE OF POISONING PLEASE PHONE / IN GEVAL VAN VERGIFTIGING SKAKEL;

0861 555777 / 082 446 8946

**Batch No:**

**Lotnommer:**

**DATE OF MANUFACTURE:**

**VERVAARDIGINGSDATUM:**

**UN No. / VN Nr. 3082 ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (Glyphosate (isopropylamine salt))**

# ERASE 360 SL

Reg. No. L6206, Act No. 36 of 1947  
Botswana Reg. No. W130314, Act No. 18 of 1999  
Malawi Reg. No. MW/PCB/2008/0120  
Namibian Reg. No. N-AR 1200

**READ THE LABEL BEFORE USE**  
**KEEP OUT OF REACH OF CHILDREN AND ANIMALS**

A NON-SELECTIVE SYSTEMIC POST-EMERGENCE HERBICIDE FOR THE CONTROL OF PERENNIAL AND ANNUAL WEEDS IN AGRICULTURE AND INDUSTRIAL AREAS.

GROUP/GROEP

9

HERBICIDE/ONKRUIDDODER

## ACTIVE INGREDIENT:

Glyphosate (Isopropylamine salt) .....480 g/l

Glyphosate (acid equivalent).....360 g /l



**DANGER**

## REGISTRATION HOLDER:

**Nulandis® (A division of AECI Limited)**

Reg. No. 1924/002590/06

P.O. Box 14418

Witfield, 1467

Tel : (011) 823 8000

## WARNINGS:

### GHS Hazard Statements

Causes skin irritations.

Causes serious eye damage.

Toxic to aquatic life with long lasting effects.

Poisonous if swallowed.

Can be corrosive to zinc and other metal equipment.

Do not mix, store or apply **ERASE 360 SL** solutions in galvanised or unlined steel (except stainless steel) containers or spray tanks.

**ERASE 360 SL** may react with such containers to form a highly combustible and explosive gas mixture.

Store in a cool, dry, well-ventilated place, away from food, feed, seed, fertilizers and other agricultural remedies.

Keep out reach of children, uninformed persons and animals.

Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing. In case of poisoning – CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM.

Aerial application: Notify all the inhabitants of the immediate vicinity of the area to be sprayed and issue the necessary warnings. Do not spray over or allow drift to contaminate water or adjacent areas. **ERASE 360 SL** is a highly active herbicide that in small quantities, when used incorrectly can cause serious damage to crop seedlings, deciduous fruit trees and grape vines during the budding – and early season growth stages.

Under the following conditions it can cause serious damage as far as 3 to 5 km from the nearest spray path of the aircraft: Cloudy weather with relative humidity above 80 % and low air movement of less than 5 km per hour. When such conditions prevail, aerial application should **NOT** be carried out where crop seedlings, deciduous fruit trees and grape vines in budding or early development stages are present within 5 km of the nearest spray path of the aircraft.

**ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL CLIMATIC AND STORAGE CONDITIONS; QUALITY OF DILUTION WATER; COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF WEEDS AGAINST THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS, WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE REGISTRATION HOLDER IN THE EVENT OF ANY UNCERTAINTY.**

### **PRECAUTIONS:**

#### **GHS Precautionary Statements**

Do not breathe dust, fume, gas, mist, vapours or spray.

Wash thoroughly after handling.

Wear protective gloves, protective clothing, eye protection & face protection

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

IF SWALLOWED: Call a POISON CENTRE or doctor/ physician if you feel unwell.

In case of accidental contact with skin wash immediately with plenty of water.

Do not eat, drink or smoke whilst mixing or applying or before washing hands and face and changing clothes.

Prevent contamination of food, feed, drinking water and eating utensils.

Prevent spray drift onto other crops, grazing or any other areas not under treatment as this will cause serious crop damage.

Clean applicator thoroughly before re-using and dispose of wash water where it will not contaminate crops, grazing, rivers or dams.

**ERASE 360 SL** is actively absorbed through immature bark and may result in localised or translocated injury to plants.

**CONTACT WITH SPRAY DRIFT OR DIRECT CONTACT WITH LEAVES OR IMMATURE BARK AND FRUIT OF DESIRED PLANTS, MUST BE AVOIDED AT ALL COST.**

Triple rinse the empty container: Invert the empty container over the spray or mixing tank and drain for at least 30 seconds after the flow has slowed down to dripping. Thereafter rinse the empty container three times in succession with one quarter of the container volume fresh water and decant the rinsate into the spray or mixing tank. Puncture the triple rinsed container and dispose of via an approved collector or recycler ([www.croplife.co.za](http://www.croplife.co.za)). Do not bury, burn or donate the container to any other parties that may use it as a container for food or beverages.

**RELEVANT SUBSTANCES:**

| Substance name<br>(IUPAC/EC)         | Concentration<br>% by weight | Classification<br>EC1272/2008                                            |
|--------------------------------------|------------------------------|--------------------------------------------------------------------------|
| Glyphosate<br>(isopropylamine salt)  | 60-70%                       | Serious eye damage (Category 1) H318.<br>Aquatic chronic 2 H411.         |
| Surfactant Blend of<br>Agrochemicals | 10%                          | Skin irritation (Category 2) H315. Eye<br>irritation (Category 2B) H320. |

**NOTE:** The other ingredients not included above, do not cause or contribute towards the correct GHS classification of **ERASE 360 SL** and are therefore, in terms of the South African Regulations for Hazardous Chemical Agents - 2021; Regulation 14(b), not listed.

**FIRST AID:**

**In case of inhalation:** If inhaled, remove to fresh air. If not breathing, give artificial respiration. If breathing is difficult, give oxygen. Get medical attention.

**In case of skin contact:** In case of contact, immediately flush skin with plenty of water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse. Thoroughly clean shoes before reuse. Get medical attention immediately.

**In case of eye contact:** Check for and remove any contact lenses. In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Cold water may be used. Get medical attention immediately.

**In case of ingestion:** Immediately transfer patient to nearest hospital or medical centre, warning by telephone of the estimated arrival time so that the start of treatment is not delayed. Do NOT induce vomiting unless directed to do so by medical personnel. Never give anything by mouth to an unconscious person. Loosen tight clothing such as a collar, tie, belt or waistband.

**NOTE TO PHYSICIAN:**

Treat symptomatically.

**RESISTANCE WARNING:**

For resistance management, **ERASE 360 SL** is a group code 9 herbicide. Any weed population may contain individuals naturally resistant to **ERASE 360 SL** and other group code 9 herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. The resistant weeds may not be controlled by **ERASE 360 SL** or any other group code 9 herbicides.

**The following weed species/weeds have been found to be resistant to glyphosate in some areas: *Conyza* spp. (Fleabane), *Lolium* spp. (Ryegrass), *Plantago lanceolata* (Narrow-leaved ribwort)**

To delay herbicide resistance:

- Avoid exclusive repeated use of herbicides from the same herbicide group code. Alternate or tank mix with registered products from different herbicide group codes.
- Integrate other control methods (chemical, cultural, biological) into weed control programmes.

For specific information on resistance management contact the registration holder of this product

**MODE OF ACTION:**

The mode of action of Glyphosate is by inhibition of 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS), an enzyme of the aromatic acid biosynthetic pathway.

## **DIRECTIONS FOR USE:** USE ONLY AS DIRECTED

### **Surfactants:**

For consistent optimum results, **ERASE 360 SL** should not be diluted beyond 1,5% in the final spray volume. Should this dilution fall below 1,5% add 500 ml per 100 l of a suitable surfactant.

### **USE RESTRICTIONS:**

**ERASE 360 SL** is a non-selective systemic herbicide, which is only active when applied post-emergence to the foliage and immature (green) bark of plants.

**ERASE 360 SL's** action is slow and the maximum herbicidal effect is only visible a few weeks after application.

Apply when weeds are actively growing in spring and summer or autumn to difficult to control perennial weeds when they are actively translocating nutrients to roots, bulbs, rhizomes and stolins.

Regrowth should be resprayed as recommended to prevent re-infestation of the treated area.

Irrigation of weeds a few days prior to treatment will ensure that weeds are actively growing and ensure maximum herbicidal efficacy.

Do not apply to wet weeds.

Do not apply to dormant weeds or weeds which suffer from moisture or temperature stress.

Rain and irrigation within 6 hours of application may reduce the efficacy.

Do not apply to weeds, which are covered with a layer of dust as efficacy will be impaired.

Pesticide run-off from treated trees onto **ERASE 360 SL** treated weeds on the orchard floor may adversely affect herbicidal efficacy.

Do not spray pesticides within 12 hours after **ERASE 360 SL** application.

**ERASE 360 SL** has no pre-emergence herbicidal activity and repeated applications are necessary to control weeds originating from seed after the previous application.

Ensure that the target weeds are fully exposed to the **ERASE 360 SL** spray.

In heterogeneous weed populations consisting of annuals and perennials, mow or spray annuals, and respray after re-growth of perennials.

Pre-plant weed control sprays on sandy soils (<10 % clay) should take place at least 7 days before sowing or transplanting any seedlings (tomato and tobacco seedlings are the most sensitive). Observe longer waiting periods on soils containing less than 10 % clay.

### **COMPATIBILITY:**

**ERASE 360 SL** is compatible with **ASC** (Reg. No. L9065), **BLADBUFF 5** for countries outside of South Africa (Reg. No. L3351), **BLADBUFF 7** (Reg. No. L8121), **IMIBOOST** (Reg. No. L7470), **LI700** (Reg. No. L6550) and **STOMAWET** (Reg. No. L6692).

For ground application only: **ERASE 360 SL** can be tank mixed with MCPA (potassium salt) and simazine with **ASC** (Reg. No. L9065,) and/or both **IMIBOOST** (Reg. No. L7470). The addition of ammonium sulphate (2%) to the spray water before adding MCPA or simazine in tank mixture with **ERASE 360 SL** improves compatibility.

However, since the formulation of other products may change without the knowledge of Nulandis and the quality of water may vary from farm to farm, the compatibility test must always be carried out prior to application to confirm compatibility.

It is recommended that a pre-test on the physical compatibility and plant safety be done before the use of such a combination with **ERASE 360 SL** in a tank mixture.

### **APPLICATION METHOD:**

#### **GROUND APPLICATION:**

**ERASE 360 SL** can be applied in spray volumes ranging from 30 - 600 l / ha.

Various types of application equipment are suitable for application viz. tractor mounted boom sprayers, knapsack sprayer, CDA applications, wiper equipment and aerial application.

Always ensure that equipment is thoroughly cleaned.

Always use high quality clean water with a low /free of organic matter and low salt content. Avoid the use of brack or muddy water, or water with a high colloid content derived from soils high in organic matter.

Calibrate the application equipment correctly under field conditions.

Do not spray when wind speeds exceeds 10 km per hour.

Ensure a fine (**NOT** a mist spray), even droplet distribution and thorough coverage of the target weeds.

Where drift may be a hazard use low pressures of 100 to 200 kPa or use low drift nozzles.

Ensure thorough coverage of the target weeds without spraying to run-off.

## AERIAL APPLICATION :

Aerial application may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 10118 (Aerial Application of Agricultural Remedies). It is important to ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria are met:

The use of a suitable drift retardant adjuvant and / or low drift nozzles (e.g. straight stream nozzles) is recommended. In the case of fixed-wing aircraft flying at a speed faster than 130 mph, the maximum deflection angle of the nozzles or spray stream, as measured from a horizontal straight backwards orientation, may not exceed 30 degrees. In the case of slower flying fixed wing aircraft the maximum deflection angle, as described above, may not exceed 55 degrees.

### Equipment:

Use suitable atomising equipment (hydraulic nozzles or rotary atomisers) that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product through drift. The operator must use a setup that will produce a droplet spectrum with the lowest possible relative span.

All nozzles / atomisers should be positioned within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.

### Application parameters:

A minimum volume of 30 – 50 l spray mixture per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aerially at a lower volume rate than recommended above.

A droplet coverage of 30-40 droplets per cm<sup>2</sup> must be recovered at the target.

A droplet spectrum with a VMD (Volume Median Diameter) of 300 - 350 microns is recommended. Ensure that the production of fine droplets (smaller than 150 microns) is restricted to a minimum.

The height of the spray boom should be maintained at 3-4 metres above the target. Do not spray when aircraft is in a climb, at the top or during a dive, or when banking.

Apply before the crop becomes too dense, thus preventing proper coverage of the weeds.

### Meteorological conditions:

The difference between the wet and dry bulb readings as determined by a whirling hygrometer, must not exceed 8 °C

Do not spray under turbulent, unstable conditions during the heat of the day when rising thermals and downdraughts occur.

Do not spray under temperature inversion conditions (spraying in or above the inversion layer).

Do not spray if the wind speed exceeds 15 km/h.

Stop spraying under turbulent, unstable and dry conditions during the heat of the day.

Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:

- reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage).
- damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.

### General:

Ensure that fields are accurately marked and that the aerial spray operator knows exactly which fields to spray.

Obtain an assurance from the aerial spray operator that the above requirements will be met.

Drift can be controlled by:

1. Addition of drift control agents.
2. The use of low drift spray nozzles.

**UNDESIRED TREES AND WOODY PLANTS**  
**ALIEN INVADERS AND DECLARED INVADING SPECIES**  
 Apply in 200 ℓ water per ha.

| <b>SPECIES CONTROLLED</b>                                                                                        | <b>ℓ / ha</b> | <b>HANDGUN<br/>ℓ / 100 ℓ<br/>water</b> | <b>KNAPSACK<br/>mℓ / 20 ℓ<br/>calibrated<br/>at 200 ℓ / ha</b> | <b>REMARKS</b>                                                                                                                                                                                                                |
|------------------------------------------------------------------------------------------------------------------|---------------|----------------------------------------|----------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <i>Acacia mearnsii</i><br>Black wattle                                                                           | 3             | 1,5                                    | 300 mℓ                                                         | Summer / autumn application.<br>Spray up to 2 m high.                                                                                                                                                                         |
| <i>Acacia saligna</i><br>Port Jackson willow<br>a) Bipinnate leaf stage<br>b) Up to 60 cm high                   | 2<br>4        | 1,5<br>1,5                             | 300 mℓ<br>300 mℓ                                               | Apply early summer to autumn.                                                                                                                                                                                                 |
| <i>Eucalyptus grandis</i><br>Blue gum<br>a) Single stem trees<br>b) Multi-stem trees                             |               | 5<br>7                                 | 1000 mℓ<br>1400 mℓ                                             | Apply as a stump treatment, after felling and removal of saw dust on same day as felling, to cambium using low pressure and a hollow cone nozzle. Should coppicing occur ringbark to expose entire cambium area and re-apply. |
| <i>Chromolaena odorata</i><br>Triffid Weed = Paraffin Weed                                                       |               | 1                                      | 200 mℓ                                                         | Apply in summer / autumn to young plants to a height of 0,5 to 1,2m. Coppices from slashed plants may require a follow-up treatment.                                                                                          |
| <i>Caesalpinea decapetala</i><br>Mauritius thorn                                                                 | 3             | 1,5                                    | 300 mℓ                                                         |                                                                                                                                                                                                                               |
| <i>Lantana camara</i><br>Lantana                                                                                 | 6             | 3                                      | 600 mℓ                                                         | Apply during summer / autumn and wet plants completely.                                                                                                                                                                       |
| <i>Solanum mauritianum</i><br>Bugtree = Bug weed<br>a) Seedlings<br><br>b) Large trees                           | <br><br>2     | 500 mℓ<br><br>1,5                      | 300 mℓ                                                         | Apply summer / autumn.<br>Spray up to a height of 1,0 m.<br><br>Apply to 1 m high coppice growth after reducing stems to ± 100 mm.                                                                                            |
| <i>Sesbania punicea</i><br>Red Sesbania<br>a) Seedlings up to 1 m<br><br>over 1 m<br><br>b) Shrubs or tall trees | 3             | 1,5<br><br>2,0<br><br>1,5 - 2          | 300 mℓ<br><br>400 mℓ<br><br>300 - 400 mℓ                       | Slash and treat coppice to height of 1 m.                                                                                                                                                                                     |

# PERENNIAL GRASSES AND BROADLEAVED WEEDS

APPLY IN 200 ℓ WATER PER ha

| SPECIES CONTROLLED                                                                                                                               | BOOM SPRAYER<br>ℓ / ha | HANDGUN<br>ℓ / 100 ℓ<br>water | KNAPSACK<br>ml / 20 ℓ<br>calibrated<br>at 200 ℓ / ha | REMARKS                                                                                                                            |
|--------------------------------------------------------------------------------------------------------------------------------------------------|------------------------|-------------------------------|------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------|
| <i>Cynodon dactylon</i><br>Common Couch =<br>Common Quick Grass<br>a) initial treatment<br>b) follow up treatment                                | 8<br>6                 | 4<br>3                        | 800 ml<br>600 ml                                     | Apply in summer and repeat in autumn or treat in autumn and follow up in summer.                                                   |
| <i>Cyperus esculentus</i><br>Yellow Nutsedge =<br>Yellow Watergrass<br><br>And<br><i>Cyperus rotundus</i><br>Purple Nutsedge = Red<br>watergrass | 6                      |                               | 600 ml                                               | Apply in summer at pre-flowering stage. If re-growth occurs, spray with 1,5 % solution or 3.0 ℓ / ha. (Best results in Feb/March). |
| <i>Eragrostis curvula</i><br>Weeping Love Grass                                                                                                  | 2                      | 1,5                           | 300 ml                                               | Apply to fully-grown grass but before onset of dormancy.                                                                           |
| <i>Paspalum dilatatum</i><br>Common Paspalum                                                                                                     | 6                      | 3                             | 600 ml                                               | Apply at flowering and re-apply to re-growth in autumn at 4 ℓ / ha.                                                                |
| <i>Paspalum paspaloides</i><br>=> <i>Paspalum distichum</i><br>Couch Paspalum =<br>Water Couch                                                   | 8                      |                               | 800 ml                                               | Apply during flowering but before seed shed. Repeat application on re-growth at 4 ℓ / ha or 300 ml / 20 ℓ.                         |
| <i>Pennisetum clandestinum</i><br>Kikuya grass<br>a) initial application<br>b) repeat application                                                | 4<br>3                 | 1,5<br>1,5                    | 400 ml<br>300 ml                                     | Apply when grass is growing vigorously and treat any re-growth to prevent re-infestation.                                          |
| <i>Panicum maximum</i><br>Ubabe = Guineagrass                                                                                                    | 6                      | 3                             | 600 ml                                               | Repeat application at 3 ℓ / ha or 1,5 % solution.                                                                                  |
| <i>Setaria megaphylla</i><br>Broadleaved Bristle<br>grass                                                                                        | 6                      | 3                             | 600 ml                                               | Apply during summer or autumn.                                                                                                     |
| <i>Plantago lanceolata</i><br>Narrow-leaved Ribwort                                                                                              | 3                      | 1,5                           | 300 ml                                               | Apply before flowering.                                                                                                            |
| <i>Malva parviflora</i><br>Small mallow                                                                                                          | 3                      |                               | 300 ml                                               | Apply before flowering.<br>Repeat after regrowth.                                                                                  |



**CONTROL OF WEEDS IN AGRICULTURE (PRE-PLANTING/REDUCED TILLAGE) AND IN  
FRUIT AND PLANTATION CROPS AND FORESTRY**

| <b>WEEDS</b>    | <b>BOOM<br/>SPRAYER<br/>ℓ / ha</b> | <b>KNAPSACK<br/>ml / 20 ℓ<br/>calibrated<br/>at 200 ℓ / ha</b> | <b>REMARKS</b>                                                                                                |
|-----------------|------------------------------------|----------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|
| Annual weeds    | 2 - 3                              | 200 - 300 ml                                                   | Apply during the 6 to 8-leaf stage. Thereafter use the higher rate on <i>Conyza</i> and <i>Commelina</i> spp. |
| Perennial weeds | -                                  | -                                                              | Refer to specific weed species as indicated in the table.                                                     |

**WEED CONTROL IN SUGARCANE**

| <b>PROBLEM</b>                                                | <b>BOOM<br/>SPRAYER<br/>ℓ / ha</b> | <b>KNAPSACK<br/>ml / 20 ℓ<br/>calibrated<br/>at 200 ℓ / ha</b> | <b>REMARKS</b>                                                                                                                                                                                      |
|---------------------------------------------------------------|------------------------------------|----------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Annual weeds                                                  | 2 – 3                              | 200 - 300 ml                                                   | Sugarcane is extremely susceptible to damage, especially during the young growing stages. Apply pre-plant overall or as a directed spray and avoid drift to prevent injury to plant or ratoon cane. |
| Perennial weeds<br><i>Cyperus rotundus</i><br>Purple nutsedge | 6                                  | 600 ml                                                         | Apply as a directed spray at early flowering and a second application at maximum re-emergence.                                                                                                      |
| Cane eradication<br>1. without undercutting roots             | 8 - 10                             | 800 – 1 000 ml                                                 | Apply at a height of 450 mm when tillering from last ratoon is complete.                                                                                                                            |
| 2. with undercutting roots                                    | 4 - 8                              | 400 - 800 ml                                                   | Apply when tillers are 350 to 1 000 mm tall in last ratoon. After 2 - 10 days undercut roots at a depth of 100 - 150 mm using a subsurface shear.                                                   |
| Diseased or off-type stools                                   |                                    | 2 ℓ                                                            | Apply as a spot treatment                                                                                                                                                                           |

### CONTROL OF WEEDS IN INDUSTRIAL AREAS

**ERASE 360 SL** can be used to control annual and perennial unwanted plants by post-emergence application on industrial sites such as roadsides, power substations, railroads etc. Consult the weed tables for specific weeds on this label

| WEEDS           | BOOM<br>SPRAYER<br>ℓ / ha | KNAPSACK<br>mℓ / 20 ℓ<br>calibrated<br>at 300 ℓ / ha | REMARKS                                                                                                                                          |
|-----------------|---------------------------|------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|
| Annual weeds    | 2 – 4                     | 133 - 266 mℓ                                         | Use the lower rate during the 6 to 8-leaf stage and the higher dosage up to seed formation. Weeds must be growing actively for maximum efficacy. |
| Perennial weeds | 4 - 10                    | 266 - 666 mℓ                                         | Apply between summer and autumn and repeat on re-growth. Protect wanted trees / plants against contact.                                          |

### CROP USE RESTRICTIONS

| CROP                                                                    | REMARKS                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|-------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>GRAPES</b>                                                           | Do not use in grapes younger than 2 years or near interplants younger than 2 years. Apply only as a directed spray from bud to dormancy. Use only during the dormant pre-budburst stage for destruction of cover crops in late winter in the Western Cape. Use only during pre-budburst between low growing bush or trellised vines or vines lower than 60 cm. Avoid spray drift contact with buds, foliage, green stems, suckers and fresh wounds. Allow a safety period of at least 10 days between pruning and application to prevent uptake from wounds. |
| <b>POME FRUIT:</b><br>Apples and Pears                                  | Use only as a directed spray in orchards older than 3 years or near interplants older than 3 years provided that green bark and foliage are protected from spray drift.                                                                                                                                                                                                                                                                                                                                                                                      |
| <b>STONE FRUIT:</b><br>Peaches, Apricots, Plums.                        | Prevent spray and spray drift contact with leaves, green stems and fruit.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>SUB TROPICAL FRUIT:</b><br>Citrus, Nuts, Avocados, Litchis, Papayas. | Prevent spray and spray drift contact with leaves, green stems and fruit. Allow a 10 day safety period after pruning before application, to prevent uptake from wounds.                                                                                                                                                                                                                                                                                                                                                                                      |

**BLADBUFF 5®**, and **BLADBUFF® 7** are the registered trademarks of AECI® Limited  
**LI 700®** is the registered trademark of LOVELAND INDUSTRIES, INC., GREELEY, USA  
**ASC**, **IMIBOOST** and **STOMAWET** are the registered tradenames of AECI® Limited