

FRUITBOOST® PHASE 1

FERTILIZER GROUP 2

A liquid organic nutrient complex for use as a foliar spray on fruit in the flowering stage to prevent or correct deficiencies.

ACTIVE INGREDIENTS:

Potassium (K) 75 g/kg (93.5 g/l)
Zinc (Zn) 7 930 mg/kg (9 888.7 mg/l)
Boron (B) 400 mg/kg (498.8 mg/l)
(Contains seaweed)
SG. 1.247 @ 20°C

NETT VOLUME:

Batch No:

Date of Manufacture:

UN No.: Not regulated



REGISTRATION HOLDER:

Nulandis® (A division of AECL Limited)

Reg. No. 1924/002590/06

P.O. Box 14418

Witfield, 1467

Tel : (011) 823 8000

www.nulandis.com

FRUITBOOST® is the registered trademark of AECL Limited (Nulandis® a division of AECL Limited)

APPLICATION TABLE:

FRUITBOOST PHASE 1 is specifically designed to be applied as a foliar spray to fruit trees during flower bud swell to full flowering, unless stated otherwise.

CROP	DOSAGE RATE	APPLICATION DETAILS
Avocados Litchis Macadamias Mangoes	500 ml / 100 l water to a maximum of 10 l/ha	Apply twice during flower bud swell to full flowering, in a minimum of 1 000 l water.
Nectarines Plums Peaches Apricots	500 ml to 1 l / 100 l water to a maximum of 5 l/ha	Apply twice during flower bud swell to full flowering.
Citrus	500 ml / 100 l water to a maximum of 10 l/ha	Apply three times during flower bud swell to full flowering, in a minimum of 1 000 l water.
Grapes	350 ml to 500 ml / 100 l water to a maximum of 10 l/ha	Apply three times during flower bud swell to full flowering. Alternate Fruitboost Phase 1 applications with Calcimax sprays.
Apples Pears	500 ml to 1 l / 100 l water to a maximum of 5 l/ha	Apply twice during flower bud swell to full flowering.

CALCIMAX® is the registered trademark of AECL Limited (Nulandis® a division of AECL Limited), Reg. No. K5999, Act No. 36 of 1947