

Fruitboost Phase 1

ZEMA Product No:

A liquid organic nutrient complex for use as a foliar spray on fruit in the flowering stage to prevent or correct deficiencies.

FERTILIZER GROUP 2

Composition:

Potassium (K) 75 g/kg (93.5 g/l)
Zinc (Zn) 7 930 mg/kg (9 888.7 mg/l)
Boron (B) 400 mg/kg (498.8 mg/l)
(Contains seaweed
SG. 1.247 @ 20 °C

Hazard Statement:

Harmful if swallowed

Manufacturer & Supplier:

Nulandis, a division of AECI LTD
14 Field Road, Lilianton, Boksburg, 1459,
South-Africa
Tel: +27 11 823 8000



Nulandis® and Qwemikelp® are registered trademarks of AECI Limited (Nulandis®, a division of AECI Limited)

Net Volume:

UN No. Not regulated

Precautionary Statements:

P264 Wash thoroughly after handling.
P270 Do not eat, drink when using this product.

Local Distributor:

ATS Agrochemicals Ltd.
Plot No 397A, Makeni,
Kafue Road, Lusaka,
Zambia.
Tel: +260 1 246 724-6



P7536

WARNING



APPLICATION TABLE:

FRUITBOOST PHASE 1 is specifically designed to be applied as a foliar spray to fruit trees during flower bud swell to full flowering, unless stated otherwise.

CROP	DOSAGE RATE	APPLICATION DETAILS
AVOCADOS LITCHIS MACADAMIAS MANGOES	500 ml / 100 ℓ water to a maximum of 10 ℓ/ha	Apply twice during flower bud swell to full flowering, in a minimum of 1 000 ℓ water.
NECTARINES PLUMS PEACHES APRICOTS	500 ml to 1 ℓ / 100 ℓ water to a maximum of 5 ℓ/ha	Apply twice during flower bud swell to full flowering.
CITRUS	500 ml / 100 ℓ water to a maximum of 10 ℓ/ha	Apply three times during flower bud swell to full flowering, in a minimum of 1 000 ℓ water.
GRAPES	350 ml to 500 ml / 100 ℓ water to a maximum of 10 ℓ/ha	Apply three times during flower bud swell to full flowering. Alternate Fruitboost Phase 1 applications with Calcimax sprays.
APPLES PEARS	500 ml to 1 ℓ / 100 ℓ water to a maximum of 5 ℓ/ha	Apply twice during flower bud swell to full flowering.

CALCIMAX® is the registered trademark of AECI Limited (Nulandis® a division of AECI Limited), Reg. No. K5999, Act No. 36 of 1947

P7538