

Fruitboost Phase 2

A liquid nutrient complex for use as a foliar spray on fruit trees in the fruit set and development stage as a foliar application to prevent or correct deficiencies.

Inorganic Foliar Fertilizer

Composition:

Calcium (Ca)	20 g/kg (24.7 g/l)
Potassium (K)	51 g/kg (62.9 g/l)
Zinc (Zn)	4 000 mg/kg (4 940 mg/l)
Boron (B)	1 500 mg/kg (1 852.5 mg/l)

(Contains seaweed)

SG: 1.235 @ 20 °C

pH: 5 - 8

DIRECTIONS FOR SAFE DISPOSAL:

Invert empty container over the spray tank and allow draining for at least 30 sec after the flow has slowed down to a drip. Thereafter rinse empty container 3 times with a volume of water equal to at least 10% of that of the container and add the rinses to the contents of the spray tank before disposing of the container. Destroy empty container and do not use for any other purpose.

SUPPLIER:

NULANDIS, a division of AECI Limited
14 Field Road
Lilianton, Boksburg, 1459, South-Africa
Tel: +27 11 823 8000
www.nulandis.com



DISTRIBUTOR:

LACHLAN
Sunflag Industrial Park, Mombasa road
P.O Box 49470, 00100 Nairobi Kenya
Telephone +254 20 207 3912/3/4
www.lachlanafrica.com



Country of origin: South-Africa

Net Volume:

Batch No.:

Manufacturing date:

Expiry date: 3 years from date of manufacture

UN No.: Not Regulated

STORAGE INSTRUCTIONS:

Store product in a segregated and approved area.
Keep container in a cool, well-ventilated area. Keep container tightly closed and sealed until ready for use.



P7548

FRUITBOOST PHASE 2

A liquid nutrient complex for use as a foliar spray on fruit trees in the fruit set and development stage as a foliar application to prevent or correct deficiencies.

Composition:

Calcium (Ca)	20 g/kg (24.7 g/l)
Potassium (K)	51 g/kg (62.9 g/l)
Zinc (Zn)	4 000 mg/kg (4 940 mg/l)
Boron (B)	1 500 mg/kg (1 852.5 mg/l)
(Contains seaweed)	

SUPPLIER:

NULANDIS , a division of AECI Limited
14 Field Road
Lilianton, Boksburg, 1459, South-Africa
Tel: +27 11 823 8000
www.nulandis.com



DISTRIBUTOR:

LACHLAN
Sunflag Industrial Park, Mombasa road
P.O Box 49470, 00100 Nairobi Kenya
Telephone +254 20 207 3912/3/4
www.lachlanafrica.com



APPLICATION TABLE:

FRUITBOOST PHASE 2 is specifically designed to be applied as a foliar spray to fruit trees during fruit set and development, unless stated otherwise.

SHAKE WELL BEFORE USE.

CROP	DOSAGE RATE	APPLICATION DETAILS
AVOCADOS LITCHIS MACADAMIAS MANGOES	500 ml / 100 l water to a maximum of 10 l/ha	Apply three times during fruit set and development, in a minimum of 1 000 l water.
PASSION FRUIT	500 ml to 1 l / 100 l water to a maximum of 5 l/ha	Apply three times during fruit set and development.
CITRUS	500 ml / 100 l water to a maximum of 10 l/ha	Apply twice during fruit set and development, in a minimum of 1 000 l water.