

Genie[®] Gro

FERTILIZER GROUP 3

Reg. No. M300, Act No. 36 of 1947

GENIE GRO is a liquid nutrient containing fulvic acid and seaweed to increase root and canopy biomass and a marketable yield.

COMPOSITION:

Nitrogen (N) 51 g/kg (54.57 g/l)
Fulvic acid 12 g/kg (12.84 g/l)

(Contains *Ecklonia maxima*)

SG 1.07 @ 20 °C

Genie Gro pH: 3.9

BATCH NO:

UN No. Not regulated

DIRECTIONS FOR USE:

Before applying **Genie Gro** consult a qualified Agronomist or your AECl Plant Health Crop Adviser on the necessity of using it.

CONTENTS: 1; 5; 10 ; 20; 25; 200; 1 000 L



Registration Holder:

AECl[®] Limited

Reg. No. 1924/002590/06

P.O. Box 14418

Witfield, 1467

Tel : (011) 823 8000

14 Field Road (Cnr Bird Street, Lilianton, 1459) Tel: (011) 823-8000

www.aeciph.com

www.aeciworld.com/

Genie[®] Gro is the registered trademark of AECl[®] Limited

WARNINGS:

Harmful when swallowed.

Do not store **Genie Gro** in direct sunlight. Do not store **Genie Gro** in temperatures below 5°C or above 45 °C.

Keep out of reach of children.

DIRECTIONS FOR USE: ONLY USE AS DIRECTED

COMPATIBILITY:

Genie Gro is compatible with **Rappid**® (Reg. No. K9404, Act. No. 36 of 1947), **Calcimax**® (Reg. No. K5999, Act. No. 36 of 1947), **Magmax**® (Reg. No. B2754, Act. No. 36 of 1947), **Manmax**® (Reg. No. B2768, Act. No. 36 of 1947) and **Zincmax**® (Reg. No. B4772, Act. No. 36 of 1947).

It is recommended to do a compatibility test on a small area and observe the treated area for 3 days before doing commercial applications. Its compatibility with other products may be influenced by several factors. As certain factors may vary from farm to farm, a physical compatibility test must always be performed before such a tank mixture is sprayed. For more information on compatibility or in the event of uncertainty, contact your local AECI Plant Health Crop Adviser.

MIXING INSTRUCTIONS:

- Fill the spray tank with ¼ of its volume with water.
- Add the required amount of **Genie Gro**.
- Fill the remainder of the spray tank with water and maintain agitation.

APPLICATION METHOD:

Apply through overhead irrigation, boom sprayer, aerial application or fertigation.

DO NOT apply **Genie Gro** when conditions are extremely hot. Best applied in temperatures in the ranges of 5 - 25°C.

CROP	TYPE OF APPLICATION	RATE / HA	RECOMMENDATION
SMALL GRAIN including Wheat (under irrigation), Barley, Oats and Rye)	Foliar	5 to 10 ℓ	Apply twice. Start the 1st application at the change from vegetative to reproductive stage (2 nd internode extending). The 2 nd application at early grain fill (end of flowering). NOTE: Do not exceed a concentration of 2% in the spray solution.

CONDITIONS OF SALE: These goods are manufactured and tested to specific standards and are sold in good faith, however no guarantee is given as conditions and methods under which they are used are beyond the control of the manufacturer. Before using, user should determine the suitability of the product for his/her intended use and user assumes the risk and liability in connection therewith.

Before applying **Genie Gro**, consult a qualified Agronomist or your local AECI Plant Health Crop Adviser on the necessity of using it.

IMPORTANT: BEFORE APPLYING GENIE GRO, CONSULT A QUALIFIED AGRONOMIST OR CROP ADVISER ON THE USE THEREOF.

Rappid[®], Calcimax[®], Magmax[®], Manmax[®] and Zincmax[®] are the registered trademarks of AECI[®] Limited

Genie[®] Gro

MISSTOF GROEP 3

Reg. Nr. M300, Wet Nr. 36 van 1947

GENIE GRO is 'n vloeibare voeding wat fulviensuur en seewier bevat, om wortel en bogrondse blaardak biomassa en 'n bemarkbare opbrengs te verhoog.

SAMESTELLING:

Stikstof (N) 51 g/kg (54.57 g/l)
Fulviensuur..... 12 g/kg (12.84 g/l)

(Bevat *Ecklonia maxima*)

SG 1.07 @ 20 °C

Genie Gro pH: 3.9

LOTNOMMER:

VN Nr. Nie gereguleer nie

GEBRUIKSAANWYSINGS:

Vóór toediening van **Genie Gro** raadpleeg 'n gekwalifiseerde Agronoom of u AECI Plant Health Gewasadviseur oor die noodsaaklikheid vir die gebruik daarvan.

INHOUD: 1; 5; 10 ; 20; 25; 200; 1 000 L



Registrasiehouer:

AECI[®] Beperk

Reg. Nr. 1924/002590/06

Posbus 14418

Witfield, 1467

Tel : (011) 823 8000

Fieldweg 14 (H/v Birdstraat, Lilianton, 1459) Tel: (011) 823-8000

www.aeciph.com

www.aeciworld.com/

Genie[®] Gro is die geregistreerde handelsmerk van AECI[®] Beperk

WAARSKUWINGS:

Skadelik indien ingesluk.

Moenie Genie Gro in direkte sonlig stoor nie. **Moenie Genie Gro** in temperature onder 5°C of bo 45 °C stoor nie.

Hou buite bereik van kinders.

GEBRUIKSAANBEVELINGS: GEBRUIK SLEGS SOOS AANGEDUI

VERENIGBAARHEID:

Genie Gro is verenigbaar met **Rappid**[®] (Reg. Nr. K9404, Wet. Nr. 36 van 1947), **Calcimax**[®] (Reg. Nr. K5999, Wet. Nr. 36 van 1947), **Magmax**[®] (Reg. Nr. B2754, Wet. Nr. 36 van 1947), **Manmax**[®] (Reg. Nr. B2768, Wet. Nr. 36 van 1947) en **Zincmax**[®] (Reg. Nr. B4772, Wet. Nr. 36 van 1947).

Dit word egter aanbeveel dat 'n toets op 'n klein area gedoen en gemonitor word vir 3 dae, voordat enige kommersiële toedienings gedoen word.

Die verenigbaarheid met ander produkte kan deur verskeie faktore beïnvloed word. Omdat hierdie faktore van tyd tot tyd kan varieër moet 'n fisiese mengbaarheidstoets eers uitgevoer word, voordat enige tenkmengsel gespuit word. Vir verdere inligting rakende verenigbaarheid of in die geval van onsekerheid, kontak u plaaslike AECI Plant Health Gewasadviseur.

VERMENGINGSINSTRUKSIES:

- Vul die spuitnek tot ¼ van die totale volume met water.
- Voeg die benodigde hoeveelheid **Genie Gro** by.
- Vul die res van die spuitnek met water en hou aan om voortdurend te roer.

TOEDIENINGSMETODE:

Dien toe deur oorhoofse besproeiing, balkbespuiting, lugtoediening of bemesting.

MOENIE toedien wanneer toestande uiters warm is nie.

Die beste tyd van toediening van **Genie Gro** is by temperature wat wissel van 5 tot 25 °C.

GEWAS	TIPE TOEDIENING	TOEDIENINGSHOEVEELHEID / HA	AANBEVELING
KLEINGRAAN insluitend Koring (onder besproeiing), Gars, Hawer en Rog)	Blaar	5 tot 10 ℓ	Dien tweekeer toe. Begin die 1ste toediening tydens die verandering vanaf vegetatiewe na reproductiewe stadium (2 ^{de} internode uitbreiding). Die 2 ^{de} toediening tydens vroeë graanvul (einde van blom). NOTA: Moenie 'n konsentrasie van 2% in die spuitoplossing oorskry nie.

VERKOOP VOORWAARDES:

Hierdie goedere word vervaardig en getoets wat aan spesifieke standarde voldoen en word in goedertrou verkoop, maar aangesien die voorwaardes en metodes waaronder hulle gebruik word buite die beheer van die vervaardiger is, word geen waarborg gegee nie. Vóór gebruik, moet die gebruiker die geskiktheid van hierdie produk vir sy/haar voorgenome gebruik bepaal.

Raadpleeg 'n gekwalifiseerde Agronoom of u AECI Plant Health Gewasadviseur, voordat **Genie Gro** toegedien word, oor die noodsaaklikheid vir die gebruik daarvan

BELANGRIK: RAADPLEEG 'N GEKWALIFISEERDE AGRONOOM OF GEWASADVISEUR VOORDAT GENIE GRO TOEGEDIEN WORD, OOR DIE NOODSAAKLIKHEID VAN DIE GEBRUIK DAARVAN.

Rappid[®], Calcimax[®], Magmax[®], Manmax[®] en Zincmax[®] is die geregistreerde handelsmerke van AECI[®] Beperk