

IMIBOOST

Reg. No./Nr L7470 , Act No./Wet Nr. 36 of/van 1947
Namibian Reg. No. / Namibiese Reg. Nr. N-AR 1667

AN ADJUVANT TO IMPROVE THE EFFICACY OF THE HERBICIDES LISTED.

‘N BEVORDERINGSMIDDEL OM DIE EFFEKTIVITEIT VAN DIE GENOEMDE ONKRUIDDODERS TE VERHOOG.

Active ingredient / Aktiewe bestanddeel :

Ammonium sulphate

.....500 g/l

Ammoniumsulfaat

Netto Volume



Net Volume



REGISTRATION HOLDER / REGISTRASIEHOUER:

Nulandis® (A division of AECI Limited) / Nulandis® (‘n Afdeling van AECI Beperk)

Reg. No. / Nr. 1924/002590/06

P.O. Box / Posbus 14418

Witfield, 1467

Tel : (011) 823 8000

IN CASE OF POISONING PLEASE PHONE / IN GEVAL VAN VERGIFTIGING SKAKEL 0861 555777/082 446 8946

Batch No:

Lotnommer:

Date of manufacture:

Datum van Vervaardiging:

UN No./VN Nr.: Not regulated / nie gereguleer nie

BLUE BAND



Pamphlet

IMIBOOST

Reg. No. L7470, Act No. 36 of 1947
Namibian Reg. No. N-AR 1667

AN ADJUVANT TO IMPROVE THE EFFICACY OF THE HERBICIDES LISTED.

Active ingredient :

Ammonium sulphate

.....500 g/l

REGISTRATION HOLDER :

Nulandis® (A division of AECI Limited)

Reg. No. 1924/002590/06

P.O. Box 14418

Witfield, 1467

Tel : (011) 823 8000

CAUTION

WARNINGS:

Handle with care.

Irritating to eyes and respiratory system.

Full cognisance must be taken of all warnings, precautions and directions for use mentioned on the labels of the products with which **IMIBOOST** will be used.

Store in a cool place away from food, feeds and seed.

Keep out of reach of children, uninformed persons and animals.

Re-entry: Do not enter treated area until spray deposit has dried unless wearing protective clothing.

In case of poisoning – CALL A DOCTOR AND MAKE THIS LABEL AVAILABLE TO HIM.

ALTHOUGH THIS ADJUVANT HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL CLIMATIC AND STORAGE CONDITIONS; QUALITY OF DILUTION WATER, COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE ADJUVANT CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS:

Do not inhale fumes or spray mist.

Wash with soap and water after use and accidental skin contact.

Do not eat, drink or smoke whilst mixing or applying, or before washing hands and face and changing clothes.

IMIBOOST may be corrosive to metal parts of sprayers.

Prevent contamination of food, feedstuffs, drinking water and eating utensils.

Thoroughly wash and rinse spray equipment after use and dispose of wash water where it will not contaminate crops, grazing, rivers and dams.

When empty, invert the container over the spray tank or mixing tank and allow it to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to at least one tenth of that of the container and add the rinsings to the contents of the spray tank. Destroy the container by perforation and flattening and dispose of it in a safe way. Do not use the container for any other purpose.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

IMIBOOST is an aqueous solution that is suitable for all spray waters and spray equipment.

IMIBOOST complexes certain dissolved salts such as Calcium and Magnesium.

IMIBOOST improves the performance of the imidazolinone herbicides.

IMIBOOST is always tank mixed with the imidazolinone herbicides and applied as a post emergence application to weeds.

Use **IMIBOOST** at a rate of 2 % (volume / volume) of the spray mixture.

Ensure thorough coverage and wetting of the weeds, especially when crop foliage is dense.

The pH and hardness of the spray water may have a market effect on the performance of the glyphosate and sulphosate containing herbicides.

MIXING INSTRUCTIONS:

Fill the spray tank three-quarters with clean water. Measure the required volume of **IMIBOOST** and pre-mix this with at least 10 ℓ water before adding to the spray tank. The other product(s) to be mixed with **IMIBOOST** should be pre-mixed in a similar fashion. Agitate the water in the spray tank and then first add the **IMIBOOST** followed by the other product(s). Fill the spray tank with water to the required level while maintaining agitation to ensure thorough mixing of the spray mixture before spraying commences. Maintain agitation while spraying. The prepared spray mixture must not be left in the spray tank for any length of time, e.g. overnight.

All spray applications must be made with suitable equipment that is in good working order and correctly calibrated to give the desired coverage for that particular method of application.

RATE OF APPLICATION:

Mix 2 ℓ **IMIBOOST** per 100 ℓ water (2%)

IMIBOOST

Reg. Nr. L7470, Wet Nr. 36 van 1947
Namibiese Reg. Nr. N-AR 1667

'N BEVORDERINGSMIDDEL OM DIE EFFEKTIVITEIT VAN DIE GENOEMDE ONKRUIDDODERS TE VERHOOG.

Aktiewe bestanddeel :

Ammoniumsulfaat500 g/l

REGISTRASIEHOUER:

Nulandis® ('n Afdeling van AECI Beperk)

Reg. Nr. 1924/002590/06

Posbus 14418

Witfield, 1467

Tel : (011) 823 8000

VERSIGTIG

WAARSKUWINGS:

Hanteer versigtig.

Irriterend vir oë en die asemhalingssisteem.

Uitvoering moet gegee word aan alle waarskuwings, voorsorgmaatreëls en gebruiksaanwysings soos aangedui op die etikette van die produkte waarmee **IMIBOOST** gebruik gaan word.

Berg in 'n koel plek weg van voedsel, voer en saad.

Hou buite bereik van kinders, oningeligte persone en diere.

Herbetreding: Moenie behandelde gebied betree voor spuitneerslag droog is nie, tensy beskermende oorklere gedra word. In geval van vergiftiging - ONTBIED 'N GENEESHEER EN STEL HIERDIE ETIKET AAN HOM BESKIKBAAR.

ALHOEWEL HIERDIE BEVORDERINGSMIDDEL OMVATTEND ONDER 'N GROOT VERSKEIDENHEID TOESTANDE GETOETS IS, WAARBORG DIE REGISTRASIEHOUER NIE DAT DIT ONDER ALLE TOESTANDE DOELTREFFEND SAL WEES NIE AANGESIEN DIE WERKING EN EFFEK DAARVAN BEÏNVLOED KAN WORD DEUR FAKTORE SOOS ABNORMALE KLIMAATS- EN BERGINGSTOESTANDE; KWALITEIT VAN VERDUNNINGSWATER, VERENIGBAARHEID MET ANDER STOWWE WAT NIE OP DIE ETIKET AANGEDUI IS NIE SOWEL AS DIE METODE, TYD EN AKKURAATHEID VAN TOEDIENING. VERDER AANVAAR DIE REGISTRASIEHOUER NIE VERANTWOORDELIKHEID VAN SKADE AAN GEWASSE, DIE OMGEWING OF VIR DIE NADELIGE EFFEK OP MENS OF DIER OF VIR 'N GEBREK AAN PRESTASIE AS GEVOLG VAN DIE VERSUIM VAN DIE GEBRUIKER OM ETIKETAANWYSINGS NA TE KOM OF AS GEVOLG VAN DIE ONTSTAAN VAN TOESTANDE WAT NIE KRAGTENS DIE REGISTRASIE VOORSIEN KON WORD NIE. RAADPLEEG DIE VERSKAFFER IN DIE GEVAL VAN ENIGE ONSEKERHEID.

VOORSORGMAATREËLS:

Moenie dampe of sproeinewel inasem nie.

Was met seep en water na gebruik en toevallige velkontak.

Moenie eet, rook of drink tydens vermenging of toediening of voordat hande en gesig gewas en skoon klere aangetrek is nie.

IMIBOOST mag invretend wees op die metaaldele van spuite.

Verhoed besoedeling van voedsel, voer, drinkwater en eetgerei.

Maak die toediener deeglik skoon voordat ander middels daarmee toegedien word en doen weg met die waswater sodat dit nie gewasse, weiding, riviere of damme sal besoedel nie.

Sodra die houer leeg is keer dit om oor die spuittenk of mengbak en dreineer vir minstens 30 sekondes nadat die vloei tot 'n gedrup verminder het. Spoel die houer daarna driekeer uit met 'n volume water gelyk aan minstens 'n tiende van die houer en gooi die spoelwater by die inhoud van die spuittenk. Vernietig die houer deur gate in te slaan en plate te druk. Moenie die houer vir enige ander doel gebruik nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

IMIBOOST is 'n wateroplosbare middel wat saam met enige spuitwater en met enige spuittoerusing gebruik kan word.

IMIBOOST komplekseer sekere oplosende soute soos Kalsium en Magnesium.

IMIBOOST verbeter die werking van imidasolinoon onkruidodders.

IMIBOOST word altyd in tenkmengsels saam met imidasolinoon onkruidodders gebruik as 'n na-opkom behandeling. Gebruik **IMIBOOST** teen 'n dosis van 2 % (volume / volume), van die spuitmengsel.

Verseker goeie bedekking en benatting van die onkruide, veral wanneer die gewas dig groei.

Waterkwaliteit (pH en soute) het 'n beduidende effek op onkruidodders (veral glifosaat en verwante onkruidodders).

VERMENGINSTRUKSIES:

Maak die spuittenk driekwart vol met skoon water. Meet die benodigde hoeveelheid **IMIBOOST** af en meng dit vooraf met minstens 10 ℓ water voordat dit by die tenk gevoeg word. Ander produk(te) wat met **IMIBOOST** gemeng word moet op 'n soortgelyke wyse eers vooraf gemeng word. Roer die water in die spuittenk en voeg dan die **IMIBOOST** eerste by gevolg deur die ander produk(te). Vul die spuittenk met water tot by die verlangde vlak terwyl daar voortdurend geroer word om deeglike vermenging te verseker alvorens daar met die toediening begin word. Handhaaf roering gedurende toediening. Aangemaakte spuitmengsels moenie in die spuittenk gelaat word vir 'n onbepaalde tyd, bv. oornag nie.

Alle toedienings moet gemaak word met geskikte toerusting wat in 'n goeie werkende toestand en korrek gekalibreer is, ten einde die verlangde bedekking te gee vir die betrokke toedieningsmetode.

TOEDIENINGSHOEVEELHEID:

Meng 2 ℓ **IMIBOOST** per 100 ℓ water (2%).