

MAGMAX



LIQUID MAGNESIUM COMPLEX

A liquid water soluble magnesium complex for use as a foliar nutrient on various crops for the prevention or correction of magnesium deficiency.

COMPOSITION:

Magnesium (Mg)..... 5.8 %w/w (7.5 %w/v)
Boron (B).....0.5 %w/w (0.65 %w/v)

pH: 5-8

DIRECTION FOR SAFE DISPOSAL:

Invert empty container over the spray tank and allow draining for at least 30 sec after the flow has slowed down to a drip. Thereafter rinse empty container 3 times with a volume of water equal to at least 10% of that of the container and add the rinses to the contents of the spray tank before disposing of the container. Destroy empty container and do not use for any other purpose.

STORAGE INSTRUCTIONS:

Store product in a segregated and approved area. Keep container in a cool, well-ventilated area. Keep container tightly closed and sealed until ready for use.

UN No: Not Regulated

Inorganic Foliar Fertiliser

SUPPLIER:

AECI Plant Health

A division of AECI Ltd
14 Field Road
Lilienton, Boksburg 1459
South-Africa
T: +27 11 823 8000
www.aeciworld.com

DISTRIBUTOR:

LACHLAN

Sunflag Industrial Park,
Mombasa Road
P.O. Box 49470, 00100
Nairobi, Kenya
T: +254 20 207 3912/3/4
www.lachlanafrica.com

Lachlan
A Unifert Company

Country of origin: South-Africa

Batch No:

Manufacturing date:

Expiry date:

Net Volume:

P7807 KEN07/21 V1

MAGMAX

A liquid water soluble organic magnesium complex for use as a foliar nutrient on various crops for the prevention or correction of magnesium deficiency.

ACTIVE INGREDIENTS:

Magnesium (Mg) 5.8 %w/w (7.5 %w/v)
Boron (B)..... 0.5 %w/w (0.65 %w/v)

MANUFACTURER & SUPPLIER:

AECI Plant Health, A division of AECI Ltd
14 Field Road
Lilianton, Boksburg 1459, South-Africa
T: +27 11 823 8000
www.aeciworld.com



DISTRIBUTOR:

LACHLAN
Sunflag Industrial Park, Mombasa Road
P.O. Box 49470, 00100, Nairobi, Kenya
T: +254 20 207 3912/3/4
www.lachlanafrica.com



DIRECTIONS FOR USE: Use only as directed

MAGMAX is compatible with most commonly used pesticides and foliar feeds, except foliar feeds with a high phosphorous content. Avoid mixing MAGMAX with highly alkaline materials such as Lime Sulphur or Bordeaux Mixture. Where overhead systems are in use, avoid scheduling irrigation directly after application. In orchards equipped with micro irrigation systems spray deposits should not be affected by irrigation. Use the lower concentration for maintenance, and the higher concentration for sufficient uptake. Apply MAGMAX as a foliar spray where neutral, rapid acting magnesium source is needed to supplement magnesium requirements.

CROP	DOSAGE ℓ /ha OR AS INDICATED	REMARKS
FRUIT CROPS Apples, Pears, Peaches, Plums	250 - 500 ml/100 ℓ water to a maximum of 5 ℓ /ha	Spray when deficiency symptoms appear. Repeat if necessary. Apply in a minimum of 1 000 ℓ water/ha.
Citrus	250 - 500 ml/100 ℓ water to a maximum of 7 ℓ /ha	Apply as a high volume spray in 1 000 ℓ water/ha. Repeat 1-2 times if necessary.
Grapes	250 - 500 ml/100 ℓ water to a maximum of 5 ℓ /ha	Spray two weeks after budding and repeat with 14 day intervals. Apply in a minimum of 1 000 ℓ water/ha.
FIELD CROPS Cereals, Peas, Canola, Maize	250 - 500 ml/100 ℓ water to a maximum of 3,5 ℓ /ha	Spray when deficiency symptoms appear. Repeat if necessary. Apply in a minimum of 300 - 500 ℓ water/ha.
HORTICULTURAL CROPS Tomatoes, Eggplants, Cucumbers, Peppers, Lettuce, Cucurbits	250 - 500 ml/100 ℓ water to a maximum of 5 ℓ /ha	Spray when deficiency symptoms appear. Repeat 3-4 times if necessary. Apply in a minimum of 200 - 400 ℓ water/ha.
Onions	250 - 500 ml/100 ℓ water to a maximum of 3,5 ℓ /ha	Apply in a minimum of 500 ℓ water/ha. Repeat 3-4 times if necessary.
Lawns	250 - 500 ml/100 ℓ water to a maximum of 3 ℓ /ha	Spray at start of growing season and repeat with 21 day intervals. Apply as a full cover spray in 450 ℓ water/ha.
Tobacco	250 - 500 ml/100 ℓ water to a maximum of 3 ℓ /ha	Apply 3-4 times with 10 day intervals in 500 ℓ water/ha.
Cotton	250 - 500 ml/100 ℓ water to a maximum of 3 ℓ /ha	Apply as soon as deficiency symptoms appear. Apply in a minimum of 300-500 ℓ water / ha. Repeat 1-2 times if necessary.
Potatoes	250 - 500 ml/100 ℓ water to a maximum of 3 ℓ /ha	Apply repeatedly in a minimum of 300 - 500 ℓ water/ha as soon as deficiency symptoms appear.
Tea	250 - 500 ml/100 ℓ water to a maximum of 7 ℓ /ha	Apply as a ground spray as soon as deficiency symptoms appear in 200 - 500 ℓ water/ha, and as an aerial application in 30-45 ℓ water/ha.

NOTE:

SPRAY VOLUMES AND DOSAGE APPLIED WILL BE DETERMINED BY THE ACTUAL TREE OR PLANT SIZE. THE DOSAGE IS DEPENDANT ON THE MAGNITUDE OF THE DEFICIENCY.

IMPORTANT:

BEFORE APPLYING MAGMAX CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURALIST ON THE NECESSITY OF USING IT.