

# MANMAX

## LIQUID MANGANESE COMPLEX FERTILIZER GROUP 2

A liquid water soluble manganese complex for use as a foliar nutrient on various crops for the prevention or correction of manganese deficiencies.

### ACTIVE INGREDIENTS:

Manganese (Mn) ..... 75 g/ℓ  
Boron (B) ..... 6.5 g/ℓ

### MANUFACTURED BY:

AECI Limited  
Reg No. 1954/003213/07  
PO Box 14418  
WITFIELD  
1467

### DIRECTIONS FOR USE: Use only as directed

**MANMAX** is compatible with most commonly used pesticides and foliar feeds, except foliar feeds with a high phosphorous content. Do not mix **MANMAX** with highly alkaline materials such as Lime Sulphur or Bordeaux Mixture. Where overhead systems are in use, avoid scheduling irrigation directly after application. In orchards equipped with micro irrigation systems spray deposits should not be affected by irrigation. Use the lower concentration for maintenance, and the higher concentration for sufficient uptake.

| CROP                                                                                                                     | DOSAGE ℓ / ha OR AS INDICATED                   | APPLICATION DETAILS                                                                                                                                                            |
|--------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Apples, Pears                                                                                                            | 75 - 100ml / 100ℓ water to a maximum of 2ℓ / ha | Apply as a foliar spray before blossom. Repeat after harvest. Apply in a minimum of 1 000 - 3 000 ℓ water/ha.                                                                  |
| Apricots, Peaches, Plums                                                                                                 | 75 - 100ml / 100ℓ water to a maximum of 2ℓ / ha | Apply as a foliar spray after harvest. Apply in a minimum of 1 000 - 3 000 ℓ water/ha.                                                                                         |
| Citrus                                                                                                                   | 75 - 100ml / 100ℓ water to a maximum of 2ℓ / ha | Apply as a foliar spray on the new flush of growth in spring. Repeat 3 - 4 weeks later if necessary. Apply in a minimum of 1 000 - 3 000 ℓ water/ha.                           |
| FIELD CROPS<br>Maize, Peas, Soyabeans, Groundnuts, Wheat, Barley, Oats, Crucifers, Onions, Tomatoes, Potatoes, Cucurbits | 75 - 100ml / 100ℓ water to a maximum of 4ℓ / ha | Apply early in the growing season when crop has sufficient leaves to absorb the spray. Repeat 3 - 4 weeks later if necessary. Apply in a minimum of 250 ℓ water/ha.            |
| Dry Beans                                                                                                                | 75 - 100ml / 100ℓ water to a maximum of 4ℓ / ha | Commence application early in the growing season when crop has sufficient leaves to absorb the spray. Repeat 2 weeks later if necessary. Apply in a minimum of 250 ℓ water/ha. |
| Olives                                                                                                                   | 75 - 100ml / 100ℓ water to a maximum of 2ℓ / ha | Apply as a foliar spray during late spring. Apply in a minimum of 1 000 - 3 000 ℓ water/ha.                                                                                    |
| Pastures                                                                                                                 | 75 - 100ml / 100ℓ water to a maximum of 7ℓ / ha | Apply in early spring and repeat annually. Apply in a minimum of 250 ℓ water/ha.                                                                                               |

### NOTE:

SPRAY VOLUMES AND DOSAGE APPLIED WILL BE DETERMINED BY THE ACTUAL TREE OR PLANT SIZE. THE DOSAGE IS DEPENDANT ON THE MAGNITUDE OF THE DEFICIENCY.

### IMPORTANT:

BEFORE APPLYING MANMAX CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURAL ADVISOR ON THE NECESSITY OF USING IT.