

MaxiBoost®

For further information see attached pamphlet

LIQUID MICRO NUTRIENT MIXTURE

FERTILIZER GROUP 2

Active ingredients / Aktiewe bestanddele:

Magnesium (Mg)	1.13 % m/v
Iron* (Fe)	0.57 % m/v
Manganese* (Mn)	0.28 % m/v
Zinc* (Zn)	0.17 % m/v
Copper* (Cu)	0.29 % m/v
Boron (B)	0.29 % m/v
Molybdenum (Mo)	0.005 % m/v
Sulfur (S)	1.46 % m/v
(Contains seaweed)	

*Chelated (EDTA)

CONTENTS / INHOUD:

BATCH NO / LOTNOMMER:

DATE OF MANUFACTURE / DATUM VAN VERVAARDIGING:

REGISTERED AND DISTRIBUTED BY:

Refiloe Trading (Pvt) Ltd
P O Box AY 201, AMBY
Bayer Complex
1 Martin Road, Msasa
Harare, Zimbabwe
Tel: +263-4-446882-8

REGISTERED AND MANUFACTURED BY:

Nulandis®
A division of AECI Ltd.
Reg. No. 1924/002590/06
P. O. Box 14418, Witfield, 1467
Tel : (011) 823 8000
www.nulandis.com

UN No. Not regulated



MaxiBoost®

LIQUID MICRO NUTRIENT MIXTURE

FERTILIZER GROUP 2

Active ingredients:

Magnesium (Mg)	1.13 % m/v
Iron* (Fe)	0.57 % m/v
Manganese* (Mn)	0.28 % m/v
Zinc* (Zn)	0.17 % m/v
Copper* (Cu)	0.29 % m/v
Boron (B)	0.29 % m/v
Molybdenum (Mo)	0.005 % m/v
Sulfur (S)	1.46 % m/v
(Contains seaweed)	

*Chelated (EDTA)

REGISTRATION HOLDER:

Nulandis® (A division of AECI Limited)
Reg. No. 1924/002590/06
P.O. Box 14418
Witfield, 1467
Tel : (011) 823 8000
www.nulandis.com



DIRECTIONS FOR USE: USE ONLY AS DIRECTED

FOLIAR APPLICATION:

Apply as a ground application in a minimum of 500 ℓ spray mixture/ha or as an aerial application in 30 ℓ spray mixture per hectare.

SOIL APPLICATION:

Apply as a drench at transplant.
Shake well before use and ensure thorough agitation during mixing and application. Do not use smaller than 50 mesh filters in the spray equipment when applying **MAXIBOOST**.

APPLICATION TABLE:

CROP	RATE	APPLICATION METHOD
BARLEY, OATS	2 ℓ/ha	Apply as a single spray at growth stage 4 - 7.
GROUNDNUTS	2 ℓ/ha	Apply the first application 2 - 3 weeks after germination and repeat 3 - 4 weeks later.
LUPINS	2 ℓ/ha	Apply a single spray 2 - 7 weeks after germination.
MAIZE	1 ℓ/ha, or 2 ℓ/ha	Apply as a row application at the 3 to 5-leaf stage. Apply as an overall spray at the 3 to 5-leaf stage with an adjuvant Bio-Film (Reg. No. L3684, Act No. 36 of 1947) or Biodew (Reg. No. L8120, Act No. 36 of 1947).
ORNAMENTALS, FLOWERS	200 mℓ/100 ℓ water	Apply as a full cover spray with 3 week intervals.
POTATOES	4 ℓ/ha 2 ℓ/ha	Apply the first spray 28 days after germination. Repeat application 10 - 14 days later, before flowering.

CROP	RATE	APPLICATION METHOD
SOYA BEANS	4 ℓ/ha 2 ℓ/ha	Apply the first spray at the third trifoliate leaf stage. Repeat application 10 - 14 days later.
TOMATOES, PEPPERS	1 mℓ / 250 mℓ water/plant 2 ℓ/ha	Apply as a drench at transplant. Repeat application as a spray at 14 day intervals for 2 sprays.
VEGETABLES: Cruciferae, Celery, Lettuce	1 mℓ / 250 mℓ water/plant 2 ℓ/ha	Apply as a drench at transplant. Repeat application as a spray at 14 day intervals for 2 sprays.
WHEAT	2 ℓ/ha	Apply at growth stage 7 and repeat at growth stage 13.
GRAPES	2 ℓ/ha 200 mℓ/100 ℓ	Apply at 2-5 cm shoot length in 500 - 1 000 ℓ water/ha. Non-bearing vines: Apply from 2-5 cm shoot length and repeat 2-3 times with 14 day intervals for root development.
TOBACCO	1 mℓ / 250 mℓ water/plant 2 ℓ/ha	Apply as a drench at transplant. Repeat application as a spray. If required, a second application can be applied 14 days later.
CITRUS	2 ℓ/ha	Apply the first application at bud break and the second application at 100% flowering.
APPLES	200 - 400 mℓ/100 ℓ to a maximum of 4 ℓ/ha	In sub-optimal fruit growing conditions, apply 4 weeks before commencing harvest and repeat 14 days later. Apply as either a high or low volume application, but ensure that the final concentration in the tank remains between 200 and 400 mℓ/100 ℓ of water.
DECIDUOUS FRUIT TREES	200 mℓ/100 ℓ	Non-bearing trees: Start applications 2-3 weeks after budbreak. Apply 2-3 applications as a foliar spray with 14 to 21 day intervals for root development. Bearing trees: Apply just after green point and repeat 14 days later.

IMPORTANT: BEFORE APPLYING MAXIBOOST CONSULT A QUALIFIED AGRONOMIST OR AN AGRICULTURAL ADVISER ON THE NECESSITY OF USING IT.

BIO-FILM® and **BIODEW®** are the registered trademarks of Nulandis® (A division of AECI Ltd.)

