

MaxiBoost®

For further information see attached pamphlet

LIQUID MICRO NUTRIENT MIXTURE

FERTILIZER GROUP 2

COMPOSITION:

Magnesium (Mg)	0.99 %m/m	(11.3 g/l)
Iron* (Fe)	0.51 %m/m	(5.7 g/l)
Manganese* (Mn)	0.25 %m/m	(2.8 g/l)
Zinc* (Zn)	0.15 %m/m	(1.7 g/l)
Copper* (Cu)	0.26 %m/m	(2.9 g/l)
Boron (B)	0.26 %m/m	(2.9 g/l)
Molybdenum (Mo)	0.005 %m/m	(0.05 g/l)
Sulfur (S)	1.3 %m/m	(14.6 g/l)

(Contains seaweed)

*Chelated (EDTA)

STORAGE AND HANDLING:

Store product in a segregated and approved area. Keep container in a cool, well-ventilated area. Keep separate from food and feed products. Keep container tightly closed and sealed until ready for use.

DIRECTIONS FOR SAFE DISPOSAL:

Invert empty container over the spray tank and allow to drain for at least 30 sec after the flow has slowed down to a drip. Thereafter rinse empty container 3 times with a volume of water equal to at least 10% of that of the container and add the rinses to the contents of the spray tank before disposing of the container. Destroy empty container and do not use for any other purpose.

CONTENTS:

BATCH NO:

DATE OF MANUFACTURE:

EXPIRY DATE:

Manufacturer:

Nulandis®

A division of AECI Ltd.

Reg. No. 1924/002590/06

c/o Field & Bird Road, Lillanton,
Boksburg, South-Africa

Tel : (011) 823 8000

www.nulandis.com

 **NULANDIS®**
unlocking nature's potential

A business of

 **AECI**
Plant & Animal
Health



Distributor:

Amiran Kenya Ltd.

Old Airport North Road, Nairobi

Tel: 0719 095000 | Fert@amirankenya.com

www.amirankenya.com

Country of origin: South - Africa
Inorganic fertilizer

P7853



MaxiBoost®

LIQUID MICRO NUTRIENT MIXTURE

FERTILIZER GROUP 2

Active ingredients:

Magnesium (Mg)	9,9 g/kg	(11,3 g/ℓ)
Iron* (Fe)	5,1 g/kg	(5,7 g/ℓ)
Manganese* (Mn)	2,5 g/kg	(2,8 g/ℓ)
Zinc* (Zn)	1,5 g/kg	(1,7 g/ℓ)
Copper* (Cu)	2,6 g/kg	(2,9 g/ℓ)
Boron (B)	2,6 g/kg	(2,9 g/ℓ)
Molybdenum (Mo)	0,05 g/kg	(0,05 g/ℓ)
Sulfur (S)	13 g/kg	(14,6 g/ℓ)
(Contains seaweed)		

*Chelated (EDTA)

REGISTRATION HOLDER:

Nulandis® (A division of AECl Limited)
Reg. No. 1924/002590/06
P.O. Box 14418
Witfield, 1467
Tel : (011) 823 8000
www.nulandis.com



08.2018 Version 04

MAXIBOOST® is the registered trademark of AECl Limited (Nulandis® a division of AECl Limited)

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

FOLIAR APPLICATION:

Apply as a ground application in a minimum of 500 ℓ spray mixture/ha or as an aerial application in 30 ℓ spray mixture per hectare.

SOIL APPLICATION:

Apply as a drench at transplant.
Shake well before use and ensure thorough agitation during mixing and application. Do not use smaller than 50 mesh filters in the spray equipment when applying **MAXIBOOST**.

APPLICATION TABLE:

CROP	RATE	APPLICATION METHOD
BARLEY, OATS	2 ℓ/ha	Apply as a single spray at growth stage 4-7.
GROUNDNUTS	2 ℓ/ha	Apply the first application 2 - 3 weeks after germination and repeat 3 - 4 weeks later.
LUPINS	2 ℓ/ha	Apply a single spray 2 - 7 weeks after germination.
MAIZE	1 ℓ/ha, or 2 ℓ/ha	Apply as a row application at the 3 to 5-leaf stage. Apply as an overall spray at the 3 to 5-leaf stage with an adjuvant Bio-Film (Reg. No. L3684, Act No. 36 of 1947) or Biodew (Reg. No. L8120, Act No. 36 of 1947).
ORNAMENTALS, FLOWERS	200 mℓ/100 ℓ water	Apply as a full cover spray with 3 week intervals.
POTATOES	4 ℓ/ha 2 ℓ/ha	Apply the first spray 28 days after germination. Repeat application 10 - 14 days later, before flowering.

CROP	RATE	APPLICATION METHOD
SOYA BEANS	4 ℓ/ha 2 ℓ/ha	Apply the first spray at the third trifoliate leaf stage. Repeat application 10 - 14 days later.
TOMATOES, PEPPERS	1 mℓ / 250 mℓ water/plant 2 ℓ/ha	Apply as a drench at transplant. Repeat application as a spray at 14 day intervals for 2 sprays.
VEGETABLES: Cruciferae, Celery, Lettuce	1 mℓ / 250 mℓ water/plant 2 ℓ/ha	Apply as a drench at transplant. Repeat application as a spray at 14 day intervals for 2 sprays.
WHEAT	2 ℓ/ha	Apply at growth stage 7 and repeat at growth stage 13.
GRAPES	2 ℓ/ha 200 mℓ/100 ℓ	Apply at 2-5 cm shoot length in 500 - 1 000 ℓ water/ha. Non-bearing vines: Apply from 2-5 cm shoot length and repeat 2-3 times with 14 day intervals for root development.
TOBACCO	1 mℓ / 250 mℓ water/plant 2 ℓ/ha	Apply as a drench at transplant. Repeat application as a spray. If required, a second application can be applied 14 days later.
CITRUS	2 ℓ/ha	Apply the first application at bud break and the second application at 100% flowering.
APPLES	200 - 400 mℓ/100 ℓ to a maximum of 4 ℓ/ha	In sub-optimal fruit growing conditions, apply 4 weeks before commencing harvest and repeat 14 days later. Apply as either a high or low volume application, but ensure that the final concentration in the tank remains between 200 and 400 mℓ/100 ℓ of water.
DECIDUOUS FRUIT TREES	200 mℓ/100 ℓ	Non-bearing trees: Start applications 2-3 weeks after budbreak. Apply 2-3 applications as a foliar spray with 14 to 21 day intervals for root development. Bearing trees: Apply just after green point and repeat 14 days later.

IMPORTANT: BEFORE APPLYING MAXIBOOST CONSULT A QUALIFIED AGRONOMIST OR AN AGRICULTURAL ADVISER ON THE NECESSITY OF USING IT.

BIO-FILM® and **BIODEW®** are the registered trademarks of Nulandis® (A division of AECl Ltd.)



P6457