

Before using this product read the label carefully

MAXIBOOST[®]

LIQUID MICRO NUTRIENT MIXTURE

FERTILIZER GROUP 2

ACTIVE INGREDIENTS:

Magnesium (Mg)	9,9 g/kg	(11.3 g/l)
Iron* (Fe)	5,1 g/kg	(5,7 g/l)
Manganese* (Mn)	2,5 g/kg	(2,8 g/l)
Zinc* (Zn)	1,5 g/kg	(1,7 g/l)
Copper* (Cu)	2,6 g/kg	(2,9 g/l)
Boron (B)	2,6 g/kg	(2,9 g/l)
Molybdenum (Mo)	0,05 g/kg	(0,05 g/l)
Sulfur (S)	13 g/kg	(14.6 g/l)
(Contains seaweed)		

*Chelated (EDTA)

Density: 1.125

CONTENTS:

BATCH NO:

DATE OF MANUFACTURE:

UN No.: Not regulated



REGISTRATION HOLDER:

Nulandis® (A division of AECI Ltd)

Reg. No. 1924/002590/06

P.O. Box 14418

Witfield, 1467

Tel : (011) 823 8000

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

FOLIAR APPLICATION:

Apply as a ground application in a minimum of 500 l spray mixture/ha or as an aerial application in 30 l spray mixture per hectare.

SOIL APPLICATION:

Apply as a drench at transplant

Shake well before use and ensure thorough agitation during mixing and application. Do not use smaller than 50 mesh filters in the spray equipment when applying **MAXIBOOST**.

APPLICATION TABLE:

CROP	RATE	APPLICATION METHOD
Barley, Oats	2 ℓ/ha	Apply as a single spray at growth stage 4 - 7.
Groundnuts	2 ℓ/ha	Apply the first application 2 - 3 weeks after germination and repeat 3 - 4 weeks later.
Lupins	2 ℓ/ha	Apply a single spray 2 - 7 weeks after germination.
Maize	1 ℓ/ha, or 2 ℓ/ha	Apply as a row application at the 3 to 5-leaf stage. Apply as an overall spray at the 3 to 5-leaf stage with an adjuvant Bio-Film (Reg. No. L3684, Act No. 36 of 1947) or BiodeW (Reg. No. L8120, Act No. 36 of 1947).
Ornamentals, Flowers	200 ml/100 ℓ water	Apply as a full cover spray with 3 week intervals.
Potatoes	4 ℓ/ha 2 ℓ/ha	Apply the first spray 28 days after germination. Repeat application 10 - 14 days later, before flowering.
Soya beans	4 ℓ/ha 2 ℓ/ha	Apply the first spray at the third trifoliate leaf stage. Repeat application 10 - 14 days later.
Tomatoes, Peppers	1 ml / 250 ml water/plant 2 ℓ/ha	Apply as a drench at transplant. Repeat application as a spray at 14 day intervals for 2 sprays.
VEGETABLES: Cruciferae, Celery, Lettuce	1 ml / 250 ml water/plant 2 ℓ/ha	Apply as a drench at transplant. Repeat application as a spray at 14 day intervals for 2 sprays.
Wheat	2 ℓ/ha	Apply at growth stage 7 and repeat at growth stage 13.
Grapes	2 ℓ/ha 200 ml/100 l	Apply at 2-5 cm shoot length in 500 - 1 000 ℓ water/ha. Non-bearing vines: Apply from 2-5 cm shoot length and repeat 2-3 times with 14 day intervals for root development.
Tobacco	1 ml / 250 ml water/plant 2 ℓ/ha	Apply as a drench at transplant. Repeat application as a spray. If required, a second application can be applied 14 days later.
Citrus	2 ℓ/ha	Apply the first application at bud break and the second application at 100% flowering
Apples	200 – 400 ml/100 l to a maximum of 4 l/ha	In sub-optimal fruit growing conditions, apply 4 weeks before commencing harvest and repeat 14 days later. Apply as either a high or low volume application, but ensure that the final concentration in the tank remains between 200 and 400 ml/100 l of water.
Deciduous fruit trees	200 ml/100 l	Non-bearing trees: Start applications 2-3 weeks after budbreak. Apply 2-3 applications as a foliar spray with 14 to 21 day intervals for root development. Bearing trees: Apply just after green point and repeat 14 days later.

IMPORTANT: BEFORE APPLYING MAXI-BOOST CONSULT A QUALIFIED AGRONOMIST OR AN AGRICULTURAL ADVISER ON THE NECESSITY OF USING IT.

BIO-FILM® and **BIODEW®** are the registered trademarks of Nulandis® (A division of AECI Ltd.)

MAXIBOOST®

VLOEIBARE MIKRO-ELEMENT MENGSEL

MISSTOF GROEP 2

AKTIEWE BESTANDDELE:

Magnesium (Mg)	9.9 g/kg	(11.3 g/l)
Yster* (Fe)	5.1 g/kg	(5.7 g/l)
Mangaan* (Mn)	2.5 g/kg	(2.8 g/l)
Sink* (Zn)	1.5 g/kg	(1.7 g/l)
Koper* (Cu)	2.6 g/kg	(2.9 g/l)
Boor (B)	2.6 g/kg	(2.9 g/l)
Molibdeen (Mo)	0.05 g/kg	(0.05 g/l)
Swael (S)	13 g/kg	(14.6 g/l)
(Bevat seewier)		

* Gecheleer (EDTA)

INHOUD:

LOTNOMMER:

DATUM VAN VERVAARDIGING:

VN Nr.: Nie gereguleer nie



REGISTRASIEHOUER:

Nulandis® ('n Afdeling van AECl Beperk)

Reg. Nr. 1924/002590/06

Posbus 14418

Witfield, 1467

Tel : (011) 823 8000

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

BLAARTOEDIENING:

Dien toe as 'n grondtoediening in 'n minimum van 500 l spuitmengsel/ha óf as 'n lugtoediening in 30 l spuitmengsel per hektaar.

GRONDTOEDIENING:

Dien toe as 'n deurdrenking tydens uitplant.

Skud goed voor gebruik en verseker deeglike roering tydens vermenging en toediening. Moenie filters kleiner as 50 maas in die spuittoerusting gebruik wanneer **MAXIBOOST** toegedien word.

TOEDIENINGSTABEL:

GEWAS	DOSIS	TOEDIENINGSMETODE
Gars, Hawer	2 ℓ/ha	Dien toe as 'n enkel bespuiting tydens groeistadium 4 tot 7.
Grondbone	2 ℓ/ha	Dien die eerste toediening 2 - 3 weke na ontkieming toe, en herhaal 3-4 weke later.
Lupiene	2 ℓ/ha	Dien toe as 'n enkel bespuiting 2 – 7 weke na ontkieming.
Mielies	1 ℓ/ha of 2 ℓ/ha	Dien toe as 'n ry-toediening tydens die 3 ^{de} tot 5-blaarstadium. Dien toe as 'n totale bespuiting tydens die 3 ^{de} tot 5-blaarstadium, tesame met 'n byvoegmiddel Bio-Film (Reg. Nr. L3684, Wet Nr. 36 van 1947) of Biodew (Reg. Nr. L8120, Wet Nr. 36 van 1947).
Sierplante, Blomme	200 ml/100 ℓ water	Dien toe as 'n voldekbespuiting, met 3 weeklikse tussenposes.
Aartappels	4 ℓ/ha 2 ℓ/ha	Dien die eerste bespuiting 28 dae na ontkieming toe. Herhaal die toediening 10-14 dae later, vóór blomstadium.
Sojabone	4 ℓ/ha 2 ℓ/ha	Dien die eerste toediening toe tydens die derde driebladblaarstadium. Herhaal die toediening 10 tot 14 dae later.
Tamaties, Soetrissies	1 ml / 250 ml water/plant 2 ℓ/ha	Dien toe as 'n deurdrenking tydens uitplant. Herhaal die toediening as 'n bespuiting met 14-dae tussenposes, vir 2 bespuitings.
GROENTE: Koolgewasse, Seldery, Blaarslaai	1 ml/250 ml water/plant 2 ℓ/ha	Dien toe as 'n deurdrenking tydens uitplant. Herhaal die toediening as 'n bespuiting met 14-dae tussenposes, vir 2 bespuitings.
Koring	2 ℓ/ha	Dien toe tydens groeistadium 7 en herhaal tydens groeistadium 13.
Druive	2 ℓ/ha 200 ml/100 ℓ	Dien toe tydens 2-5 cm lootlengte, in 500 - 1 000 ℓ water/ha. Nie-draende wingerd: Dien toe vanaf 2-5 cm lootlengte. Herhaal 2-3 keer met 14-dae tussenposes vir wortelontwikkeling.
Tabak	1 ml / 250 ml water/plant 2 ℓ/ha	Dien toe as 'n deurdrenking tydens uitplant. Herhaal die toediening as 'n bespuiting. Indien nodig kan 'n tweede bespuiting 14 dae later gedoen word.
Sitrus	2 ℓ/ha	Dien die eerste bespuiting toe tydens bot en die tweede bespuiting tydens 100% blom.

Appels	200-400 ml/100 ℓ tot 'n maksimum van 4 ℓ/ha	In sub-optimale vruggroei omstandighede, dien toe 4 weke vóór aanvang van oes en herhaal 14 dae later. Dien toe as 'n hoë óf lae volume bespuiting, maar verseker dat die finale konsentrasie in die tenk tussen 200 en 400 ml/100 ℓ water is.
Sagtevrugte bome	200 ml/100 ℓ	<u>Nie-draende bome:</u> Begin die toediening 2-3 weke na bot. Dien 2-3 keer toe as 'n blaarbespuiting met 14 tot 21-dae tussenposes vir wortelontwikkeling. <u>Draende bome:</u> Dien toe kort na groenpunt en herhaal 14 dae later.

BELANGRIK: RAADPLEEG 'N GEKWALIFISEERDE AGRONOM OF LANDBOUKUNDIGE ADVISEUR VOORDAT MAXI-BOOST TOEGEDIEN WORD, OOR DIE NOODSAAKLIKHEID VAN DIE GEBRUIK DAARVAN.

BIO-FILM® en **BIODEW®** is die geregistreerde handelsmerke van Nulandis® ('n Afdeling van AECI Beperk)