

NUTRIFEED 1

Reg. No. K7882, Act No. 36 of 1947
Namibian Reg. No. N-F 1678

FERTILIZER GROUP 1

A water soluble fertilizer for use on non-bearing pome and stone fruit trees after transplant.

COMPOSITION:

Nitrogen (N)	310 g/kg
Boron (B).....	43 g/kg
Copper*(Cu)	9 g/kg

*EDTA Chelated

Batch No:

UN No: Not regulated

CONTENTS: 500 g, 1 kg, 5 kg, 10 kg, 25 kg



REGISTRATION HOLDER:

AECI® Limited

Reg. No. 1924/002590/06

P.O. Box 14418

Witfield, 1467

14 Field Road (Cnr Bird Street) Lilianton, 1459

Tel : (011) 823 8000

www.aeciph.com/www.aeciworld.com

RECOMMENDATIONS:

For use to prevent deficiencies of Copper and Boron in non-bearing pome and stone fruit trees.

Apply **Nutrifeed 1** as a foliar spray either in the early morning or late afternoon.

Do not spray when temperatures exceed 30 °C or on wilted plants as this can result in reduced uptake.

Nutrifeed 1 - a water soluble fertilizer has a mildly acidifying effect on spray waters thus making it ideal to use. However, kindly refer to "Compatibility", below for further information.

COMPATIBILITY:

Do not mix Nutrifeed 1 with lime sulphur or Bordeaux mixtures.

Do not use a buffer or acidifying agent when applying **Nutrifeed 1**.

Nutrifeed 1 is a water soluble fertilizer and it has a mild acidifying effect on spray waters thus making it ideal for use with other products. However, the compatibility of **Nutrifeed 1** has not yet been proven with all agricultural remedies. Its compatibility with other products may be influenced by several factors. As certain factors may vary from farm to farm, a physical compatibility test must always be performed before such a tank mixture is sprayed. For more information on compatibility or in the event of uncertainty, contact your local AECI Plant Health Crop Adviser. It is recommended to do a compatibility test on a small area and observe the treated area for 3 days before doing commercial applications.

DIRECTIONS FOR USE: Use only as directed

It is recommended to use a wetting agent, e.g. **Biodew**® (Reg. No. L8120) at a rate of 25 ml/100 l water.

Apply as a high volume spray.

RATE OF APPLICATION:

Apply Nutrifeed 1 at a rate of 500 g to 1 kg / 100 l water

TIME OF APPLICATION:

- 1) Start application of **Nutrifeed 1** approximately 3 weeks after beginning of budbreak.
- 2) Follow up with **Nutrifeed 2** (Reg. No. K11077, Act No. 36 of 1947) 7 to 10 days later.
- 3) After 7 to 10 days, start again with **Nutrifeed 1** and repeat this cycle until 3 applications of each product have been made.

Biodew® is the registered trademark of AECI® Limited

Nutrifeed 2 is the registered tradename of AECI® Limited

NUTRIFEED 1

Reg. Nr. K7882, Wet Nr. 36 van 1947

MISSTOF GROEP 1

'n Wateroplosbare misstof vir gebruik op nie-draende kern- en steenvrugte bome na uitplant.

SAMESTELLING:

Stikstof (N)		310 g/kg
Boor (B)		43 g/kg
Koper* (Cu)		9 g/kg

*Gecheleerde EDTA

Lotnommer:

VN Nr. Nie gereguleer nie

INHOUD: 500 g, 1 kg, 5 kg, 10 kg, 25 kg



REGISTRASIEHOUER:

AECI® Beperk

Reg. Nr. 1924/002590/06

Posbus 14418

Witfield, 1467

Fieldweg 14 (h/v Birdstraat), Lilianton, 1459

Tel : (011) 823 8000

www.aeciph.com/ www.aeciworld.com

AANBEVELING:

Vir gebruik om tekorte van Koper en Boor in nie-draende kern- en steenvrugte bome te voorkom. Dien **Nutrifeed 1** toe as 'n blaartoediening in die vroeë môre óf laatmiddag. Moenie toedien as temperature 30 °C oorskry nie, en ook nie op verlepte plante nie, want dit kan die opname verlaag.

Nutrifeed 1 - water oplosbare misstof het 'n effense versurings effek op die spuitwater en is dus ideaal om te gebruik. Die is egter raadsaam om na "Verenigbaarheid" hieronder te verwys, vir verdere inligting.

VERENIGBAARHEID:

Moenie **Nutrifeed 1** met kalkswael of Bordeaux mengsel vermeng nie.

Moenie 'n buffer- of versuringsmiddel gebruik in gevalle waar **Nutrifeed 1** toegedien word.

Nutrifeed 1 is 'n wateroplosbare misstof met 'n effense versurings effek op die spuitwater en is dus ideaal om saam met produkte te gebruik. Nieteenstaande, is die verenigbaarheid van **Nutrifeed 1** met alle landboumiddels, nog onbekend. Die verenigbaarheid met ander produkte kan deur verskeie faktore beïnvloed word. Omdat hierdie faktore van tyd tot tyd kan varieër moet 'n fisiese mengbaarheidstoets eers uitgevoer word, voordat enige tenkmengsel gespuit word. Vir verdere inligting rakende die verenigbaarheid van die produk of in die geval van onsekerheid, kontak u plaaslike AECI Plant Health Gewasadviseur. Dit word egter aanbeveel dat 'n toets op 'n klein area gedoen en gemonitor word vir 3 dae, voordat enige kommersiële toedienings gedoen word.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui

Gebruik die benattingsmiddel **Biodew®** (Reg. Nr. L8120) teen die aanbevole toediening van 25 ml/100 ℓ water.

Dien toe teen 'n hoë volume bespuiting.

TOEDIENINGSHOEVEELHEID:

Dien Nutrifeed 1 toe teen 500 g tot 1 kg / 100 ℓ water
--

TYD VAN TOEDIENING:

- 1) Begin toediening van **Nutrifeed 1** ongeveer 3 weke na aanvang van bot.
- 2) Volg op met **Nutrifeed 2** (Reg. Nr K11077, Wet Nr 36 van 1947) 7 tot 10 dae later.
- 3) Na afloop van 7 tot 10 dae, begin weer met **Nutrifeed 1** en herhaal hierdie siklus tot 3 toedienings van elke produk gedoen is.

Biodew® is die geregistreerde handelsmerk van AECI® Beperk

Nutrifeed 2 is die geregistreerde handelsnaam van AECI® Beperk