

NUTRIFEED 2

Reg. No. K11077, Act No. 36 of 1947
Namibian Reg. No. N-F 1679

FERTILIZER GROUP 1

A water soluble fertilizer for use on non-bearing pome and stone fruit trees after transplant.

COMPOSITION:

Nitrogen	(N)	188 g/kg
Phosphorous	(P)	143 g/kg
Zinc	(Zn)	20 g/kg
Manganese	(Mn)	33 g/kg
Sulphur	(S)	29 g/kg

Batch No:

UN No: 3077 ENVIRONMENTALLY HAZARDOUS SUBSTANCE, SOLID, N.O.S
(zinc sulphate)

CONTENTS: 1 200 kg; 250 g; 500 g; 1 kg; 5 kg; 10 kg; 20 kg; 25 kg; 50 kg



REGISTRATION HOLDER:

AECI® Limited

Reg. No. 1924/002590/06
P.O. Box 14418
Witfield, 1467

14 Field Road (Cnr Bird Street),
Lilianton, 1459
Tel : (011) 823 8000
www.aeciph.com/www.aeciworld.com

RECOMMENDATIONS:

For use to stimulate growth and prevent deficiencies of phosphorous (P), zinc (Zn) and manganese (Mn), in non-bearing pome and stone fruit trees.

Apply **Nutrifeed 2** as a foliar spray either in the early morning or late afternoon. Do not spray when temperatures exceed 30 °C or onto wilted plants as this may result in reduced uptake.

COMPATIBILITY:

Do not mix Nutrifeed 2 with lime sulphur or Bordeaux mixtures.

Do not use a buffer or acidifying agent when applying **Nutrifeed 2**.

Nutrifeed 2 is a water soluble fertilizer and it has a mild acidifying effect on spray waters thus making it ideal for use with other products. However, the compatibility of **Nutrifeed 2** has not yet been proven with all agricultural remedies. Its compatibility with other products may be influenced by several factors. As certain factors may vary from farm to farm, a physical compatibility test must always be performed before such a tank mixture is sprayed. For more information on compatibility or in the event of uncertainty, contact your local Nulandis representative. It is recommended to do a compatibility test on a small area and observe the treated area for 3 days before doing commercial applications.

DIRECTIONS FOR USE: Use only as directed

Use the wetting agent Biodew (Reg. No. L8120) at a rate of 25 ml/100 l water.

Apply as a high volume spray.

APPLICATION RATE:

Apply Nutrifeed 2 at a rate of 500 g to 1 kg /100 l water
--

TIMING OF APPLICATION:

- 1) Apply **Nutrifeed 2** - 7 to 10 days after **Nutrifeed 1** (Refer to the **Nutrifeed 1** Reg. No. K7882, Act. No. 36 of 1947 label for full directions).
- 2) Repeat 7 to 10 days after each **Nutrifeed 1** (Reg. No. K7882, Act. No. 36 of 1947) application, up to 3 applications per season.

NUTRIFEED 1 - Reg. No. K7882, Act No. 36 of 1947 is the registration of AECI® Limited
BIODEW® is the registered trademark of AECI® Limited

NUTRIFEED 2

Reg. Nr. K11077, Wet Nr. 36 van 1947

MISSTOF GROEP 1

'n Wateroplosbare misstof vir gebruik op nie-draende kern- en steenvrugte bome na uitplant.

SAMESTELLING:

Stikstof	(N)	188 g/kg
Fosfor	(P)	143 g/kg
Sink	(Zn)	20 g/kg
Mangaan	(Mn)	33 g/kg
Swael	(S)	29 g/kg

Lotnommer:

VN Nr. 3077

INHOUD: 1 200 kg; 250 g; 500 g; 1 kg; 5 kg; 10 kg; 20 kg; 25 kg; 50 kg



REGISTRASIEHOUER:

AECI® Beperk

Reg. Nr. 1924/002590/06

Posbus 14418

Witfield, 1467

Fieldweg 14 (h/v Birdstraat),

Lilianton, 1459

Tel : (011) 823 8000

www.aeciph.com/www.aeciworld.com

AANBEVELING:

Vir gebruik om groei te stimuleer en tekorte van fosfor (P), sink (Zn) en mangaan (Mn) in nie-draende kern- en steenvrugte bome te voorkom.

Dien **Nutrifeed 2** toe as 'n blaartoediening in die vroeë môre óf laatmiddag. Moenie toedien as temperature 30 °C oorskry nie, en ook nie op verlepte plante nie, want dit kan die opname verlaag.

VERENIGBAARHEID:

Moenie **Nutrifeed 2** met kalkswael of Bordeaux mengsel vermeng nie.

Moenie 'n buffer- of versuringsmiddel gebruik in gevalle waar **Nutrifeed 2** toegedien word.

Nutrifeed 2 is 'n wateroplosbare misstof met 'n effense versurings effek op die spuitwater en is dus ideaal om saam met produkte te gebruik. Nieteenstaande, is die verenigbaarheid van **Nutrifeed 2** met alle landboumiddels, nog onbekend. Die verenigbaarheid met ander produkte kan deur verskeie faktore beïnvloed word. Omdat hierdie faktore van tyd tot tyd kan varieër moet 'n fisiese mengbaarheidstoets eers uitgevoer word, voordat enige tenkmengsel gespuut word. Dit word egter aanbeveel dat 'n toets op 'n klein area gedoen en gemonitor word vir 3 dae, voordat enige kommersiële toedienings gedoen word.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui

Gebruik die benattingsagent Biodew (Reg. Nr. L8120) teen 'n aanbeveling van 25 ml/100 l water.

Dien toe teen 'n hoë volume bespuiting.

TOEDIENINGSHOEVEELHEID:

Dien Nutrifeed 2 toe teen 500 g tot 1 kg / 100 l water
--

TYD VAN TOEDIENING:

- 1) Begin toediening van **Nutrifeed 2** 7 tot 10 dae na **Nutrifeed 1** (verwys na **Nutrifeed 1** Reg. Nr. K7882, Wet Nr. 36 van 1947 se etiket vir besonderhede), toegedien is.
- 2) Herhaal 7 tot 10 dae na elke **Nutrifeed 1** toediening (Reg. Nr. K7882, Wet Nr. 36 van 1947) tot 3 toedienings per seisoen gedoen is.

NUTRIFEED 1 - Reg. Nr. K7882, Wet Nr. 36 van 1947 is die registrasie van AECI® Beperk
BIODEW® is die geregistreerde handelsmerk van AECI® Beperk