

QWEMIKELP

LIQUID NUTRIENT FOLIAR FEED COMPLEX

Liquid nutrient foliar feed complex with seaweed for use on crops as indicated.

COMPOSITION:

Nitrogen (N)	7.7 %w/w
Potassium (K)	0.8 %w/w
Iron (Fe)	0.11 %w/w
Boron (B)	0.15 %w/w
Manganese (Mn)	0.05 %w/w
Molybdenum (Mo)	0.02 %w/w

(Contains seaweed)

pH: 5-8

DIRECTION FOR SAFE DISPOSAL:

Invert empty container over the spray tank and allow draining for at least 30 sec after the flow has slowed down to a drip. Thereafter rinse empty container 3 times with a volume of water equal to at least 10% of that of the container and add the rinses to the contents of the spray tank before disposing of the container. Destroy empty container and do not use for any other purpose.

UN No: Not Regulated

Inorganic Foliar Fertiliser

SUPPLIER:

AECI Plant Health
A division of AECI Ltd
14 Field Road
Lilienton, Boksburg 1459
South-Africa
T: +27 11 823 8000
www.aeciworld.com

DISTRIBUTOR:

LACHLAN
Sunflag Industrial Park,
Mombasa Road
P.O. Box 49470, 00100
Nairobi, Kenya
T: +254 20 207 3912/3/4
www.lachlanafrica.com

Lachlan
A Unifert Company

STORAGE INSTRUCTIONS:

Store product in a segregated and approved area. Keep container in a cool, well-ventilated area. Keep container tightly closed and sealed until ready for use.

Country of origin: South-Africa

Batch No:

Manufacturing date:

Expiry date:

Net Volume:

QWEMIKELP

Liquid nutrient foliar feed complex with seaweed for use on crops as indicated.

COMPOSITION:

Nitrogen (N)	7.74 %w/w	(8.73 %w/v)
Potassium (K)	0.88 %w/w	(0.99 %w/v)
Sulfur (S)	0.087 %w/w	(0.09 %w/v)
Magnesium (Mg)	0.375 %w/w	(0.423 %w/v)
Iron (Fe)	0.1125 %w/w	(0.1269 %w/v)
Boron (B)	0.1502 %w/w	(0.1694 %w/v)
Manganese (Mn)	0.0564 %w/w	(0.0636 %w/v)
Zinc (Zn)	0.0370 %w/w	(0.0417 %w/v)
Copper (Cu)	0.0200 %w/w	(0.0225 %w/v)
Molybdenum (Mo)	0.0237 %w/w	(0.0267 %w/v)

(Contains seaweed)

MANUFACTURER & SUPPLIER:

AECI Plant Health, A division of AECI Ltd
14 Field Road
Lilienton, Boksburg 1459, South-Africa
T: +27 11 823 8000
www.aeciworld.com



DISTRIBUTOR:

LACHLAN
Sunflag Industrial Park, Mombasa Road
P.O. Box 49470, 00100, Nairobi, Kenya
T: +254 20 207 3912/3/4
www.lachlanafrica.com



DIRECTIONS FOR USE: USE ONLY AS DIRECTED
Shake well before use.

Ground application:

Always dilute in a maximum 600 l water/ha and apply at a rate of 2 to 4 l/ha.

Aerial application:

Dilute in 30 – 40 l water/ha and apply at a rate of 2 to 4 l/ha.

APPLICATION TABLE:

CROP	APPLICATION METHOD
Citrus	Apply at bud swell and repeat at 21 day intervals.
Cotton	Apply at the 10-leaf stage and repeat with 14 day intervals.
Legume crops	Apply at the 4 th trifoliate and repeat with 14 day intervals.
Maize	Apply at the 5 to 7-leaf stage.
Potatoes	Apply at the 6-leaf stage and repeat 14 days later.
Sugarcane	Apply once the cane has reached a height of 60 cm.
Fruiting Vegetables: Cucumbers, Egg plant, Melons, Peppers, Squash and Tomatoes	Apply at the 6-leaf stage and repeat with 14 day intervals.
Leaf Vegetables: Asparagus, Broccoli, Brussels sprout, Cabbages, Cauliflower, Celery, Chives, Lettuce, Parsley and Spinach	Apply at the 6-leaf stage and repeat with 14 day intervals.
Root Vegetables: Beets, Carrots, Turnip, Parsnip and Radish	Apply at the 6-leaf stage and repeat with 14 day intervals.
Wheat	Apply at the 5 to 7-leaf stage.
Sunflower	Apply at the 5 to 7-leaf stage and repeat 14 days later.