

QWEMIKELP

Namibian Reg. No. N-F 1469

LIQUID NUTRIENT FOLIAR FEED COMPLEX

GROUP 2 FERTILIZER

LIQUID NUTRIENT FOLIAR FEED COMPLEX WITH SEAWEED FOR USE ON CROPS AS INDICATED.

ACTIVE INGREDIENTS:

Nitrogen (N)	77.4 g/kg	(87.3 g/l)
Phosphorus (P)	10.7 g/kg	(12 g/l)
Potassium (K)	8.8 g/kg	(9.9 g/l)
Sulfur (S)	0.87 g/kg	(0.9 g/l)
Magnesium (Mg)	3.75 g/kg	(4.23 g/l)
Iron (Fe)	1125 mg/kg	(1269 mg/l)
Boron (B)	1502 mg/kg	(1694 mg/l)
Manganese (Mn)	564 mg/kg	(636 mg/l)
Zinc (Zn)	370 mg/kg	(417 mg/l)
Copper (Cu)	200 mg/kg	(225 mg/l)
Molybdenum (Mo)	237 mg/kg	(267 mg/l)
(Contains seaweed)		

CONTENTS:

BATCH NO:

DATE OF MANUFACTURE:

UN No.: Not regulated



REGISTRATION HOLDER:

Nulandis® (A division of AECI Ltd)

Reg. No. 1924/002590/06

P.O. Box 14418

Witfield, 1467

Tel : (011) 823 8000

www.nulandis.com

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Shake well before use.

Ground application:

Always dilute in a maximum 600 ℓ water/ha and apply at a rate of 2 to 4 ℓ/ha.

Aerial application:

Dilute in 30 – 40 ℓ water/ha and apply at a rate of 2 to 4 ℓ/ha.

APPLICATION TABLE:

CROP	APPLICATION METHOD
Citrus	Apply at bud swell and repeat at 21 day intervals.
Cotton	Apply at the 10-leaf stage and repeat with 14 day intervals.
Legume crops	Apply at the 4 th trifoliolate and repeat with 14 day intervals.
Maize	Apply at the 5 to 7-leaf stage.
Potatoes	Apply at the 6-leaf stage and repeat 14 days later.
Sugarcane	Apply once the cane has reached a height of 60 cm.
Fruiting Vegetables: Cucumbers, Egg plant, Melons, Peppers, Squash and Tomatoes	Apply at the 6-leaf stage and repeat with 14 day intervals.
Leaf Vegetables: Asparagus, Broccoli, Brussels sprout, Cabbages, Cauliflower, Celery, Chives, Lettuce, Parsley and Spinach	Apply at the 6-leaf stage and repeat with 14 day intervals.
Root Vegetables: Beets, Carrots, Turnip, Parsnip and Radish	Apply at the 6-leaf stage and repeat with 14 day intervals.
Wheat	Apply at the 5 to 7-leaf stage.
Sunflower	Apply at the 5 to 7-leaf stage and repeat 14 days later.

QWEMIKELP

Namibiese Reg. Nr. N-F 1469

‘N VLOEIBARE BLAARVOEDING KOMPLEKS

MISSTOF GROEP 2

‘N VLOEIBARE BLAARVOEDING KOMPLEKS MET SEEWIER VIR GEBRUIK OP DIE
GENOEMDE GEWASSE SOOS AANGEDUI.

AKTIEWE BESTANDDELE:

Stikstof (N)	77.4 g/kg	(87.3 g/l)
Fosfaat (P)	10.7 g/kg	(12 g/l)
Kalium (K)	8.8 g/kg	(9.9 g/l)
Swael (S)	0.87 g/kg	(0.9 g/l)
Magnesium (Mg)	3.75 g/kg	(4.23 g/l)
Yster (Fe)	1125 mg/kg	(1269 mg/l)
Boor (B)	1502 mg/kg	(1694 mg/l)
Mangaan (Mn)	564 mg/kg	(636 mg/l)
Sink (Zn)	370 mg/kg	(417 mg/l)
Koper (Cu)	200 mg/kg	(225 mg/l)
Molibdeen (Mo)	237 mg/kg	(267 mg/l)
(Bevat seewier)		

INHOUD:

LOTNOMMER:

DATUM VAN VERVAARDIGING:

VN Nr: Nie gereguleer nie



REGISTRASIEHOUER:

Nulandis® (‘n Afdeling van AECI Bpk)

Reg. Nr. 1924/002590/06

Posbus 14418

Witfield, 1467

Tel : (011) 823 8000

www.nulandis.com

GEBRUIKSAANWYSINGS:

GEBRUIK SLEGS SOOS AANGEDUI

Skud goed voor gebruik.

Grondtoediening:

Verdun in 'n maksimum 600 ℓ water/ha en dien toe teen 'n dosis van 2 tot 4 ℓ/ha.

Lugtoediening:

Verdun in 30 – 40 ℓ water/ha en dien toe teen 'n dosis van 2 tot 4 ℓ/ha.

TOEDIENINGSTABEL:

GEWAS	TOEDIENINGSMETODE
Sitrus	Dien toe tydens knopswel en herhaal met 21-dae tussenposes.
Katoen	Dien toe by die 10-blaarstadium en herhaal met 14-dae tussenposes.
Peulgewasse	Dien toe by die 4 ^{de} driebladblaarstadium en herhaal met 14-dae tussenposes.
Mielies	Dien toe tydens die 5 tot 7-blaarstadium.
Aartappels	Dien toe tydens die 6-blaarstadium en herhaal 14 dae later.
Suikerriet	Dien toe sodra die suikerriet 'n hoogte van 60 cm bereik het.
Vrugdraende Groentes: Komkommers, Eiervrug, Spanspek, Soetrissies, Skorsies en Tamaties	Dien toe tydens die 6-blaarstadium en herhaal met 14-dae tussenposes.
Blaargroentes: Aspersies, Broccoli, Brusselse spruite, Kopkool, Blomkool, Seldery, Grasuie, Kropslaai, Pietersielie en Spinasie	Dien toe tydens die 6-blaarstadium en herhaal met 14-dae tussenposes.
Wortel- of Knolgroentes: Beet, Wortels, Witwortel, Raap en Radys	Dien toe tydens die 6-blaarstadium en herhaal met 14-dae tussenposes.
Koring	Dien toe tydens die 5 tot 7-blaarstadium.
Sonneblom	Dien toe tydens die 5 tot 7-blaarstadium en herhaal 14 dae later.