

# QWEMIKELP

## 7-2-1

### GUARANTEED ANALYSIS

|                                                      |        |
|------------------------------------------------------|--------|
| Total Nitrogen (N)                                   | 7.00%  |
| 6.5% Urea Nitrogen                                   |        |
| 0.3% Ammoniacal Nitrogen                             |        |
| 0.2% Nitrate Nitrogen                                |        |
| Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) | 2.00%  |
| Soluble Potash (K <sub>2</sub> O)                    | 1.00%  |
| Magnesium (Mg)                                       | 0.50%  |
| Boron (B)                                            | 0.150% |
| Iron (Fe)                                            | 0.113% |
| Manganese (Mn)                                       | 0.056% |
| Molybdenum (Mo)                                      | 0.024% |

**Derived from:** Aqua Ammonia, Urea, Potassium Nitrate, Kelp extract (*Ecklonia maxima*), Potassium Hydroxide, Magnesium Oxide, Boric Acid, Ferrous Sulphate, Manganese Oxide, Sodium Molybdate

### ALSO CONTAINS NON-PLANT FOOD

#### INGREDIENTS:

50% Kelp extract (*Ecklonia maxima*), (microbe food)

**F003086**

### GUARANTEED BY:

**Nulandis (A division of AECI® Ltd)**

AECI Chem Park, Block No 2 – 2nd Floor  
 200 Bergivier Drive, Chloorkop  
 Kempton Park 1619  
 South Africa  
 T: +27 11 823 8000  
[www.aeciworld.com](http://www.aeciworld.com)



**Net Content:** 2.5 Gallons (9.46 liters)

**Density:** 10.52 lbs/US gal @ 68° F

**Net Weight:** 26.3 lbs (11.92 kg)

**WARNING:** Product contains boron and the use of the fertilizing material on any crops other than those recommended may result in serious injury to the crop(s).

**WARNING:** The application of fertilizing material containing Molybdenum (Mo) may result in forage crops containing levels of Molybdenum (Mo) which are toxic to ruminant animals.

### KEEP OUT OF REACH OF CHILDREN



### WARNING

Causes serious eye irritation. Causes skin irritation. May cause respiratory irritation. Wear protective gloves. Wear eye or face protection. Use only outdoors or in a well-ventilated area. Avoid breathing vapor. Wash hands thoroughly after handling.

**UN No: Not regulated**

**P8036 USA07/21 V1**

**FIRST AID:**

**If in eye:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If on skin:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**If swallowed:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**DIRECTIONS FOR USE: USE ONLY AS DIRECTED INCLUDED BUT NOT LIMITED TO THE CROPS LISTED BELOW**

Shake well before use.

**Foliar application:**

Always dilute in a maximum of 60-gallons water per acre and apply at a rate of 1 to 2 quarts per acre.

**Aerial application:**

Dilute in 3 to 4 gallons water per acre and apply at a rate of 1 to 2 quarts per acre.

**In furrow application:**

0.9 to 1 quart per acre applied in 5-10 gallons per acre.

**APPLICATION TABLE:**

| <b>CROP</b>                                                                                                                                                     | <b>APPLICATION METHOD</b>                                                 |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------|
| <b>CITRUS</b>                                                                                                                                                   | Apply at bud swell and repeat at 21-day intervals.                        |
| <b>CORN</b>                                                                                                                                                     | Apply at the 5 to 7-leaf stage.                                           |
| <b>COTTON</b>                                                                                                                                                   | Apply at the 10-leaf stage and repeat with 14-day intervals.              |
| <b>HEMP</b>                                                                                                                                                     | Apply at the 5 to 7-leaf stage                                            |
| <b>LEGUME CROPS</b>                                                                                                                                             | Apply at the 4 <sup>th</sup> trifoliate and repeat with 14-day intervals. |
| <b>POTATOES</b>                                                                                                                                                 | Apply at the 6-leaf stage and repeat 14 days later.                       |
| <b>SUGARCANE</b>                                                                                                                                                | Apply once the cane has reached a height of 23 inches.                    |
| <b>FRUITING VEGETABLES:</b><br><b>CUCUMBERS, EGG<br/>PLANT, MELONS,<br/>PEPPERS, SQUASH AND<br/>TOMATOES</b>                                                    | Apply at the 6-leaf stage and repeat with 14-day intervals.               |
| <b>LEAF VEGETABLES:</b><br><b>ASPARAGUS,<br/>BROCCOLI, BRUSSELS<br/>SPROUT, CABBAGES,<br/>CAULIFLOWER, CELERY,<br/>CHIVES, LETTUCE,<br/>PARSLEY AND SPINACH</b> | Apply at the 6-leaf stage and repeat with 14-day intervals.               |
| <b>ROOT VEGETABLES:</b><br><b>BEETS, CARROTS,<br/>TURNIP, PARSNIP AND<br/>RADISH</b>                                                                            | Apply at the 6-leaf stage and repeat with 14-day intervals.               |
| <b>WHEAT</b>                                                                                                                                                    | Apply at the 5 to 7-leaf stage.                                           |
| <b>SUNFLOWER</b>                                                                                                                                                | Apply at the 5 to 7-leaf stage and repeat 14 days later.                  |

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>