

QWEMIKELP®

LIQUID NUTRIENT FOLIAR FEED COMPLEX

GROUP 2 FERTILIZER

LIQUID NUTRIENT FOLIAR FEED COMPLEX WITH SEAWEED FOR USE ON CROPS AS INDICATED.

COMPOSITION:

Nitrogen (N)	77.4 g/kg	(87.3 g/l)
Phosphorus (P)	10.7 g/kg	(12 g/l)
Potassium (K)	8.8 g/kg	(9.9 g/l)
Sulfur (S)	0.87 g/kg	(0.9 g/l)
Magnesium (Mg)	3.75 g/kg	(4.23 g/l)
Iron (Fe)	1125 mg/kg	(1269 mg/l)
Boron (B)	1502 mg/kg	(1694 mg/l)
Manganese (Mn)	564 mg/kg	(636 mg/l)
Zinc (Zn)	370 mg/kg	(417 mg/l)
Copper (Cu)	200 mg/kg	(225 mg/l)
Molybdenum (Mo)	237 mg/kg	(267 mg/l)
(Contains seaweed)		



DANGER

GHS HAZARD STATEMENTS

May damage fertility.

May damage the unborn child

GHS PRECAUTIONARY STATEMENTS

Wash thoroughly after handling.

Do not eat, drink when using this product .

MANUFACTURED BY:

AECI Ltd

14 Field Road, Lilianton,
Boksburg, 1459, South Africa

Tel +27 11 823 8000

www.aeciph.com

Qwemikelp® is a registered trademark of AECI® Limited

CONTENTS:

BATCH NO

UN No.: Not regulated

WARNINGS:

GHS HAZARD STATEMENTS	May damage fertility.
-----------------------	-----------------------

May damage the unborn child.

PRECAUTIONS:

GHS PRECAUTIONARY STATEMENTS
Prevention
Obtain, read, and follow all safety instructions before use.
Wash thoroughly after handling.
Do not eat, drink when using this product.
Wear impervious rubber gloves and boots, protective clothing, and chemical safety goggles.
Response
If exposed or concerned, get medical advice.
Storage
Store locked up. Disposal
Dispose of contents/container in accordance with local and national regulations.

RELEVANT SUBSTANCES:

Substance name (IUPAC/EC)	Concentration % by weight	Classification
Boric Acid	<1%	Reproductive Toxicity (Category 1B), H360FD.

NOTE: The other ingredients not included above, do not cause or contribute towards the correct GHS classification of QWEMIKELP® and are therefore, in terms of the South African Regulations for Hazardous Chemical Agents - 2021; Regulation 14(b), not listed.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED
Shake well before use.

Ground application:

Always dilute in a maximum 600 ℓ water/ha and apply at a rate of 2 to 4 ℓ/ha.

Aerial application:

Dilute in 30 – 40 ℓ water/ha and apply at a rate of 2 to 4 ℓ/ha.

APPLICATION TABLE:

CROP	APPLICATION METHOD
Citrus	Apply at bud swell and repeat at 21 day intervals.
Cotton	Apply at the 10-leaf stage and repeat within 14 day intervals.
Legume crops	Apply at the 4 th trifoliate and repeat with 14 day intervals.
Maize	Apply at the 5 to 7-leaf stage.
Potatoes	Apply at the 6-leaf stage and repeat 14 days later.
Sugarcane	Apply once the cane has reached a height of 60 cm.
Fruiting Vegetables: Cucumbers, Egg plant, Melons, Peppers, Squash and Tomatoes	Apply at the 6-leaf stage and repeat with 14 day intervals.
Leaf Vegetables: Asparagus, Broccoli, Brussels sprout, Cabbages, Cauliflower, Celery, Chives, Lettuce, Parsley and Spinach	Apply at the 6-leaf stage and repeat within 14 day intervals.
Root Vegetables: Beets, Carrots, Turnip, Parsnip and Radish	Apply at the 6-leaf stage and repeat with 14 day intervals.
Wheat	Apply at the 5 to 7-leaf stage.
Sunflower	Apply at the 5 to 7-leaf stage and repeat 14 days later.