



A business of
AECI
Plant & Animal
Health

RAPPID

9-34-0

GUARANTEED ANALYSIS

Total Nitrogen (N)9%
9% Ammoniacal Nitrogen
Available Phosphate (P_2O_5)34%


Derived from: Aqua ammonia, Ammonium polyphosphate

F003086

GUARANTEED BY:

Nulandis (A division of AECI® Ltd)
AECI Chem Park, Block No 2 - 2nd Floor
200 Bergrevier Drive, Chloorkop
Kempton Park, 1619
South Africa
T: +27 11 823 8000
www.aeciworld.com

UN No: Not regulated



Net Content: 2.5 Gallons (9.46 L)

Density: 11.43 lbs/US gal @ 68° F

Net Weight: 28.6 lbs (12.96 kg)

KEEP OUT OF REACH OF CHILDREN



WARNING

Wear protective gloves. Wear eye or face protection. Use only outdoors or in a well-ventilated area. Avoid breathing vapor. Wash hands thoroughly after handling.

P8427 USA07/21 V1

160(h) x 130(w)

CYAN

MAG

YEL

BLK

FIRST AID:

If in eye: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED – INCLUDED BUT NOT LIMITED TO THE CROPS LISTED BELOW

MIXING INSTRUCTIONS:

- Half fill tank to the required volume of water.
- Add the required amount of **RAPPID**.
- Fill the remainder of the tank with water and maintain agitation.

COMPATIBILITY:

For the purpose of compatibility, it is not recommended to mix any other agricultural products with **RAPPID**.

APPLICATION TABLE:

CROP	APPLICATION TYPE	APPLICATION RATE	WATER VOLUME	RECOMMENDATIONS
APPLES	Foliar	1.5 to 2 quarts/100 gallons of water to a maximum of 2.25 quarts RAPPID /acre	Apply in a minimum of 80 gallons/acre	Apply within one week after full bloom, up to 4 applications with a 14-day interval period.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply preferably through the irrigation system during the vegetative phase.
BEANS & CUCURBITS	Foliar	2.5 to 3 quarts/100 gallons of water to a maximum rate of 3.25 quarts RAPPID /acre	Apply in a minimum of 20 gallons of water/acre	Apply during fruit cell division phase. DO NOT apply during flowering.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply during the vegetative growth stage including the fruit cell division phase.
CANOLA & HEMP	Foliar	2.5 to 3 quarts/100 gallons of water to a maximum rate of 3.25 quarts RAPPID /acre	Apply in a minimum of 10 gallons/acre	Begin spraying from the 3-leaf stage until flowering.
	Soil	1 to 1.6 gallons/acre	Apply in 10-20 gallons/acre	Apply at the vegetative stage prior to flowering.
	Irrigation	1 to 3 gallons/acre	0.1 to 0.2 inches irrigation	Apply at plant in P-retentive soils during the vegetative stage.
	In-furrow	0.5 gallons/acre	Apply in 10-20 gallons/acre	Band place RAPPID (2 inches beneath and 2 inches away from planted seed).

CARROTS, POTATOES, TOMATOES, BRASSICA (COLE), PEPPERS, ONIONS	Foliar	1.5 to 2 quarts/100 gallons of water	Apply in a minimum of 20 gallons/acre	Apply from plant until fruit cell division phase. DO NOT apply during flowering.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply during the vegetative growth stage including the fruit cell division phase.
	In-furrow	0.5 gallons/acre	Apply in 10-20 gallons/ acre	Band place RAPPID (2 inches away from the seedlings).
CITRUS, AVOCADOS, OLIVES	Foliar	1.5 to 2 quarts/100 gallons of water and a maximum rate of 4.25 quarts RAPPID /acre	Apply in 100-320 gallons/acre	Apply during fruit cell division phase. DO NOT apply during flowering.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply during the vegetative growth phase including fruit cell division phase.
CORN	Foliar	1.5 to 2 quarts/100 gallons of water and a maximum rate of 4.25 quarts RAPPID /acre	Apply in 100-320 gallons/acre	Apply during fruit cell division phase. DO NOT apply during flowering.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply during the vegetative growth phase including fruit cell division phase.
	In-furrow	0.5 gallons/acre	Apply in 10-20 gallons/acre	Band place RAPPID (2 inches beneath and 2 inches away from planted seed).
LEGUMES, PEANUTS	Foliar	1.5 to 2 quarts/100 gallons of water	Apply in a minimum of 20 gallons/acre	Apply during the vegetative stage up to flowering.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply 3 to 4 weeks after emergence.
	In-furrow	0.5 gallons/acre	Apply in 10-20 gallons/ acre	Band place RAPPID (2 inches beneath and 2 inches away from planted seed).
PASTURES	Soil	1 to 2 gallons / acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply during Spring and Autumn flush.
	Foliar	1 quart/100 gallons of water to a maximum rate of 1.25 quarts RAPPID / acre	Apply in a minimum of 80 gallons/acre	Apply within one week after full bloom, up to 4 applications with a 14-day interval period.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply preferably through the irrigation system during the vegetative stage.

PEARS	Foliar	1 quart/100 gallons of water to a maximum rate of 1.25 quarts RAPPID /acre	Apply in a minimum of 80 gallons/acre	Apply within one week after full bloom, up to 4 applications with a 14-day interval period.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply preferably through the irrigation system during the vegetative stage.
ROOIBOS TEA	Foliar	2 to 5 quarts/100 gallons of water	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply after cutting. Use the lower dosage for small bushes and the higher dosage for large bushes.
	Soil	1 to 3 gallons/acre	Apply in a minimum of 80 gallons/acre	Apply during the vegetative stage.
STONE FRUIT	Foliar	2 to 2.5 quarts/100 gallons of water to a maximum of 2.5 quarts RAPPID /acre	Apply in a minimum of 80 gallons/acre	Apply during the vegetative stage.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply preferably through the irrigation system during the vegetative stage.
STRAWBERRIES	Foliar	1 to 1.5 quarts/100 gallons of water	Apply in a minimum of 20 gallons/acre	Apply during fruit cell division phase. DO NOT apply during flowering.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply during the vegetative growth stage including fruit cell division phase.
SUGARCANE (PLANT AND RATOON)	Foliar	0.3 to 0.5 gallons/acre	Apply in 10-20 gallons/acre	Apply the lower rate on ratoon cane when the sugarcane is 12 inches in length and the higher rate when at a height of 24 inches and when the foliar nitrogen content is low.
SUGARCANE (PLANT)	Soil	1 to 3 gallons/acre	Apply in 10-20 gallons/acre	Apply at planting in furrow.
TREE NUTS (including almonds, cashews, chestnuts, hazelnuts, pecans, pistachios, walnuts)	Foliar	3 to 3.5 quarts/100 gallons of water to a maximum of 3.5 quarts RAPPID /acre	Apply in 100-300 gallons/acre	Apply 4 weeks after full bloom. DO NOT apply during flowering.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	Apply through the irrigation system during the vegetative stage.
MACADAMIAS	Foliar	3 to 3.5 quarts/100 gallons of water to a maximum of 3.5 quarts RAPPID /acre	Apply in 100-300 gallons/acre	Apply 4 weeks after full bloom. DO NOT apply during flowering. DO NOT exceed 4 applications.
VEGETABLE SEEDLINGS (NURSERIES)	Foliar	1 to 2 quarts/100 gallons of water	Apply in a minimum of 0.02 gallons/sq	Apply 2 applications two weeks after emergence, with a 10 to 14-day interval period.

WHEAT, BARLEY, OATS, TRITICALE	Foliar	2 to 3 quarts/100 gallons of water	Apply in a minimum of 20 gallons/acre	Apply at the 4 to 6-leaf stage.
	Soil	1 to 1.6 gallons/acre	Apply in 10-20 gallons/ acre	Apply during the vegetative stage before flowering.
	Irrigation	1 to 3 gallons/acre	0.1 to 0.2 inches irrigation	Apply at plant in P-retentive soils during the vegetative stage.
	In-furrow	0.5 gallons/acre	Apply in 10-20 gallons/ acre	Band place RAPPID (2 inches beneath and 2 inches away from planted seed).
WINE AND TABLE GRAPES	Foliar	2 to 2.5 quarts/100 gallons of water to a maximum of 2.5 quarts RAPPID /acre	Apply in a minimum of 80 gallons/acre	Apply from bud burst up to two weeks prior to flowering and two weeks after flowering during the cell division phase.
	Soil	1 to 3 gallons/acre	Apply at a maximum rate of 8 gallons RAPPID /260 gallons water	

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.aapfco.org/metals.html>