

Rappid® K

POTASSIUM POLYPHOSPHATE

FERTILIZER GROUP 1

Composition:

Phosphorus (P) 74 g/kg (102.12 g/l)
Potassium (K) 132 g/kg (182.16 g/l)

SG 1.30 – 1.38 @ 20 °C
Rappid K pH 6.0 to 7.0

Contents:

1 Metric Ton; 1 000 ℓ; 210 ℓ; 25 ℓ; 20 ℓ; 5 ℓ; 1 ℓ; 500 ml

Batch no.:

UN No. Not regulated



Registration Holder:

Nulandis® (A division of AECI Limited)
Reg. No. 1924/002590/06
P.O Box 14418 Witfield, 1467
14 Field Road (Cnr Bird Street) Lilianton, 1459
Tel: (011) 823-8000
www.nulandis.com / www.aeciworld.com

Rappid® is a registered trademark of AECI® Limited

WARNINGS:

DO NOT use **Rappid K** with any calcium-containing fertilizers.

PROPERTIES:

- A potassium based product, with no nitrogen, for reproductive growth nutrition.
- **Rappid K** has a near-neutral pH of 6.0 to 7.0.
- **Rappid K** is highly plant available due to its pure, concentrated formulation. It is effective as an instant source of P and K, but also has a sustained effect on the nutrient levels of the plant.
- Ease and better control of application rates and placement of nutrients.
- Contains (P) in the polyphosphate form.

COMPATIBILITY:

Rappid K is compatible with **Alexin®** (Reg. No. B3835, Act No. 36 of 1947). For the purpose of energy enhancement it is not recommended to mix any other agricultural remedies with **Rappid K**.

MIXING INSTRUCTIONS:

- Half fill the tank to the required volume of water.
- Add the required amount of **Rappid K**.
- Fill the remainder of tank with water and maintain agitation.

APPLICATION TABLE

CROP	APPLICATION TYPE	DOSAGE	WATER VOLUME/ha	RECOMMENDATIONS
APPLES	Foliar	500 ml / 100 ℓ water and a maximum of 5 ℓ RAPPID K /ha	Apply in a minimum of 750 ℓ	Apply up to 4 applications with a 10 – 14 day interval period from 6 weeks after flowering up to 2 weeks prior to harvesting.
BEANS & CUCURBITS	Foliar	750 ml / 100 ℓ water	Apply in a minimum of 400 ℓ	Apply during fruit cell enlargement phase up to 2 weeks prior to harvesting.
CANOLA	Foliar	750 ml / 100 ℓ water	Apply in a minimum of 100 ℓ	Apply 2 applications after flowering with a 10 – 14 day interval period.
CARROTS, POTATOES, TOMATOES, BRASSICA (COLE CROPS), (PEPPERS, ONIONS)	Foliar	750 ml / 100 ℓ water	Apply in a minimum of 400 ℓ	Apply every 2 weeks after flowering until harvesting.
CITRUS, AVOCADOS, OLIVES	Foliar	750 ml / 100 ℓ water and a maximum of 10 ℓ RAPPID K /ha	Apply in 1 000 – 3 000 ℓ	Apply every 2 weeks during fruit enlargement phase until harvesting.
COTTON	Foliar	750 ml / 100 ℓ water	Apply in a minimum of 400 ℓ	Apply 10 days after flowering.
LEGUMES, GROUNDNUTS	Foliar	750 ml / 100 ℓ water	Apply in a minimum of 300 ℓ	Apply twice after flowering and podfill.
MAIZE	Foliar	750 ml / 100 ℓ water	Apply in a minimum of 300 ℓ	Apply 2 applications after pollination with a 10 – 14 day interval period.
PEARS	Foliar	500 ml / 100 ℓ water and a maximum of 5 ℓ RAPPID K /ha	Apply in a minimum of 750 ℓ	Apply up to 4 applications with a 10-14 day interval from 6 weeks after flowering until 2 weeks prior to harvesting.
ROOIBOS TEA	Foliar	500 ml / 100 ℓ water	Apply in a minimum of 100 ℓ	Apply after stress conditions.
STONE FRUIT	Foliar	500 ml / 100 ℓ water and a maximum of 5 ℓ RAPPID K /ha	Apply in a minimum of 750 ℓ	Apply up to 4 applications with a 10-14 day interval from 6 weeks after flowering until 2 weeks until harvesting.
STRAWBERRIES	Foliar	250 ml / 100 ℓ water	Apply in a minimum of 300 ℓ	Apply every 2 weeks during cell enlargement phase until harvesting.

SUGARCANE (PLANT & RATOON)	Foliar	3 to 5 ℓ /ha	Apply in a minimum of 200 ℓ	Apply the lower rate on ratoon cane when the sugarcane is 30 cm in length and the higher rate when at a height of 60 cm. Apply when the (P) and (K) levels in the leaves are low and during stress conditions.
TREE NUTS (including almonds, cashews, chestnuts, hazelnuts, pecans, pistachios, walnuts)	Foliar	<u>NON-BEARING TREES</u> 500 ml / 100 ℓ water		Apply from 6 weeks after flowering until the end of February.
MACADAMIAS		<u>BEARING TREES</u> 750 ml / 100 ℓ water and a maximum of 7.5 ℓ RAPPID K /ha	Apply in 1 000 – 3 000 ℓ	MACADAMIAS ONLY: Apply a maximum of 4 applications from 6 weeks after flowering until the end of February.
VEGETABLE SEEDLINGS (NURSERIES)	Foliar	250 ml / 100 ℓ water	Apply in a minimum of 100 ℓ	Apply every 2 weeks after transplanting.
WHEAT, BARLEY, OATS, TRITICALE	Foliar	500 ml / 100 ℓ water	Apply in a minimum of 100 ℓ	Apply from ear forming and after flowering, but not during flowering.
WINE AND TABLE GRAPES	Foliar	500 ml / 100 ℓ water and a maximum of 5 ℓ RAPPID K /ha	Apply in a minimum of 750 ℓ	Apply during cell enlargement phase up to 2 weeks prior to harvesting.

NOTE: For soil application, please consult your AECl Plant Health Crop Adviser for further assistance.

Alexin® is the registered trademark of AECl® Limited