

SOMO

FERTILIZER GROUP 2

Namibian Reg. No. N-F 1416

A foliar application or seed dressing for the prevention and correction of molybdenum deficiency.

ACTIVE INGREDIENT:

Sodium Molybdate390 g/kg

MANUFACTURED AND REGISTERED BY:

Gouws & Scheepers Pty Ltd

Reg. No. 1954/003213/07

Trading as PLAASKEM

P O BOX 14418

WITFIELD

1467



DIRECTIONS FOR USE:

CROP	RATE OF APPLICATION	REMARKS
Cucurbits	2 g/l	Let the seed soak overnight in a 0,2 % solution and plant directly afterwards.
Crucifers	100 – 250 g/ha	Apply as a foliar spray to young plants in sufficient water.
Lucerne	125 – 250 g/ha	Seed treatment simultaneously with inoculation.
Grass/Clover Pastures	155 g/ha	Foliar spray onto young re-growth after cutting.
Maize	50 g/ 1,25 l water	Place 100 kg seed in a drum with watertight lid. Add the solution and rotate the drum for 15 - 20 minutes. The seed may be treated anytime before planting.
Sunflower	25 g/ 25 kg Seed 5 g/100 l water to a maximum of 75 g/ha	A solution containing SOMO should be applied uniformly to the seed. Apply to the seedlings as a full foliar spray.
Tomatoes	250 g/ha	Apply to planting furrow before transplanting or as a foliar spray. Molybdenum deficiency occurs throughout the Transvaal Lowveld particularly on acid soils.
Soybeans	75 - 125 g/ha	Apply to the seedlings as a foliar spray.

REPEAT TREATMENT:

Depending on the quantity of SOMO applied, the following intervals between reapplication are recommended.

QUANTITY OF SOMO APPLIED PER/ha	INTERVAL PERIODS BETWEEN TREATMENTS
Less than 75 g 75 - 150 g More than 150 g	1 year 2 - 3 years 4 - 5 years

IMPORTANT:

Before applying this fertiliser, consult a qualified Agronomist or Agricultural Advisor on the necessity of using it.

SOMO

MISSTOF GROEP 2

Namibiese Reg. Nr. N-F 1416

'n Blaarbespuiting of 'n saadbehandeling om 'n molibdeen tekort te voorkom of reg te stel..

AKTIEWE BESTANDDEEL:

Natrium molibdaat 390 g/kg

VERVAARDIG EN GEREGISTREER DEUR:

Gouws & Scheepers Edms Bpk
Reg. No. 1954/003213/07
Handeldrywend as PLAASKEM
POSBUS 14418
WITFIELD
1467



GEBRUIKSAANWYSINGS:

GEWAS	DOSIS VAN TOEDIENING	OPMERKINGS
Rankgewasse	2 g/l	Week die saad oornag in 'n oplossing van 0,2 % en plant dadelik daarna.
Koolgewasse	100 – 250 g/ha	Dien toe as 'n blaarbespuiting op jong plante met voldoende water.
Lusern	125 – 250 g/ha	Behandel saad tydens ovulasie.
Grasse/Klawer Weiding	155 g/ha	Dien toe as 'n blaarbespuiting op jong hergroei na afsny.
Mielies	50 g/ 1,25 l water	Plaas 100 kg saad in 'n drom met 'n waterdigte deksel. Voeg die oplossing by en draai vir 15 - 20 minute. Die saad kan behandel word enige tyd voor plant.
Sonneblom	25 g/ 25 kg Saad 5 g/100 l tot 'n maksimum van 75 g/ha	'n Oplossing van SOMO moet eweredig op die saad aangewend word. Dien toe op saailinge as 'n blaarbespuiting.
Tamaties	250 g/ha	Dien toe in die plantry net voor aanplantings of as 'n blaarbespuiting. Molibdeen tekorte kom oor die algemeen in die Transvaal Laeveld voor veral op suurgrond.
Sojabone	75 - 125 g/ha	Dien toe op saailinge as 'n blaarbespuiting.

HERHALINGS BEHANDELING:

Afhangende van die hoeveelheid van SOMO toegedien, word die volgende tussenposes tussen toediening aanbeveel.

HOEEVEELHEID SOMO TOEGEDIEN/ha	TOEDIENINGS TUSSENPOSES
Minder as 75 g 75 - 150 g Meer as 150 g	1 jaar 2 - 3 jaar 4 - 5 jaar

BELANGRIK:

Raadpleeg 'n gekwalifiseerde Agronoom of Landboukundige voordat hierdie misstof toegedien word or die noodsaaklikheid vandie gebruik daarvan.