

CAUTION: KEEP OUT OF REACH OF CHILDREN

ToneUp™

Microbial bio-fertiliser liquid

Fertiliser group 3

Reg. No. M267, Act No. 36 of 1947

Active constituent: *Bacillus* spp 221 x 10⁹ cfu / gm

Shelf Life: 11 months

Nett volume 50ml / 500ml / 1L

SG = 1.043 at 22.3°C

ToneUp™ is a plant growth stimulant for growth improvement

Not classified as hazardous under GHS

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE OR OPENING



Microbial Plus (Pty) Ltd
Co. Reg. No. 2011/136780/07
Unit 3 Windsor Glen, Windsor Ave, Ramsgate, 4285
info@microbialplus.co.za
www.microbialplus.co.za
+27 82 906 4478

Batch No.

Manufactured date:

WARNINGS:

- Handle with care
- Do not inhale spray mix
- Harmful if swallowed
- Avoid contact with skin, eyes or inhalation and ingestion
- No cold-chain required, store at room temperature
- Store away from food, drinks and animal feed
- Keep out of reach of children, uninformed persons and animals

PRECAUTIONS:

No adverse reports have been received. However, the following precautions constitute good handling procedures and safe farming practices

- Use protective clothing, gloves and boots during mixing and application
- If skin or eye contact occurs rinse with large amounts of water
- Destroy empty packaging and containers and do not reuse for any other purpose

DIRECTIONS FOR PRODUCT PREPARATION:

- Shake well before use

When used under unfavourable soil and climate condition, or not applied according to directions, the manufacturer cannot guarantee root and soil colony formation

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

COMPATIBILITY:

Climate conditions, chemical compatibility and co-application:

- ToneUp™ can be co-applied with all known agricultural chemicals and fertilisers and under all climate conditions
- Contact the help line on +27 82 906 4478

APPLICATION METHODS:

- ToneUp™ can be applied in as much water as is required by all known methods including in-furrow application, seed application, pivot, boom spray, micros, drippers and aerial spray
- Can be applied in any temperature at any time of the day in direct sunlight

APPLICATION RATES:

CROPS	APPLICATION RATE	RECOMMENDATIONS
VEGETABLES e.g. Carrots	50ml / Ha	The application rates are based on a plant density of 100 carrots per m². Application 1: Apply at planting either as a seed application or in-furrow or as a combination of both. Application 2: At 2 weeks apply to the soil surface and water as required. Application 3: At 4 weeks repeat application to the soil surface and water as required.