

VEGEBOOST® Plus

Reg. No. B5658, Act No 36 of 1947

FERTILIZER GROUP 2

A liquid nutrient complex for foliar feeding or correction of nutritional deficiencies of vegetables.

COMPOSITION:

Calcium (Ca)	55.0 g/kg
Magnesium (Mg)	10.1 g/kg
Boron (B)	4.0 g/kg
Zinc (Zn)	2.5 g/kg
Molybdenum (Mo)	0.8 g/kg
Iron (Fe)	2.6 g/kg

SG @ 20°C 1.26

NETT MASS / NETTO MASSA

1; 5; 10; 20 & 25 kg



DANGER

GHS HAZARD STATEMENTS

Causes serious eye irritation.
May damage fertility.
May damage the unborn child.

GHS PRECAUTIONARY STATEMENTS

Obtain, read, and follow all safety instructions before use.
Do not eat, drink when using this product.



Registration Holder:

Nulandis® (A division of AECI Limited)

Reg. No.1924/002590/06

P.O Box 14418 Witfield, 1467

14 Field Road (Cnr Bird Street, Lilianton, 1459

Tel: (011) 823-8000

www.aeciph.com / www.aeciworld.com

Vegeboost® is a registered trademark of AECI® Limited

Batch no./ Lotnommer:

UN No.: **Not regulated / nie gereguleer nie**

WARNINGS:**GHS HAZARD STATEMENTS**

Causes serious eye irritation.
May damage fertility.
May damage the unborn child.

PRECAUTIONS:**GHS PRECAUTIONARY STATEMENTS****Prevention**

Obtain, read, and follow all safety instructions before use.
Do not eat, drink when using this product.
Wear impervious rubber gloves and boots, protective clothing, and chemical safety goggles.
Wash hands and face thoroughly after handling.
Do not touch eyes.

Response

If exposed or concerned, get medical advice.
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical help.

Storage

Store locked up.

Disposal

Dispose of contents/container in accordance with local and national regulations.

RELEVANT SUBSTANCES:

Substance name (IUPAC/EC)	Concentration % by weight	GHS Classification
Boric Acid	1 – 5%	Reproductive Toxicity (Category 1B), H360FD.
Ethoxylated fatty alcohol	<10%	Acute toxicity oral (Category 4) H302. Serious eye damage (Category 1) H318

NOTE: The other ingredients not included above, do not cause or contribute towards the correct GHS classification of **VEGEBOOST Plus** and are therefore, in terms of the South African Regulations for Hazardous Chemical Agents - 2021; Regulation 14(b), not listed.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

CROP	APPLICATION RATE	REMARKS
Tomatoes, Peppers, Beans, Peas, Onions	500 ml/100 ℓ water to a maximum of 4 ℓ/ha	Begin applications prior to fruit set and repeat at 7 – 10 day intervals.
Lettuce, Cauliflower, Cabbage, Chinese cabbage	500 ml/100 ℓ water to a maximum of 3 ℓ/ha	Apply weekly, ensuring good wetting of the entire plant beginning prior to onset of head formation.
Potatoes	500 ml/100 ℓ water to a maximum of 4 ℓ/ha	Apply at tuber initiation and repeat every 7 – 10 days.
Cucurbits (Pumpkins, Watermelons, Melons)	500 ml/100 ℓ water to a maximum of 4 ℓ/ha	Apply at flowering and repeat with 14 day intervals.

IMPORTANT: Before applying **VEGEBOOST PLUS** consult your local AECI Plant Health Crop Adviser on the necessity of using it.

