

ZALEXIN

READ THE LABEL CAREFULLY BEFORE OPENING
For full information see attached leaflet

A liquid organic nutrient complex containing salicylic acid.

FERTILIZER GROUP 2

ACTIVE INGREDIENT

Nitrogen (N).....	0.98 % m/v
Potassium (K).....	4.86 % m/v
Calcium (Ca).....	2.84 % m/v
Magnesium (Mg).....	0.91 % m/v
Zinc (Zn).....	2.30 % m/v
Boron (B).....	0.25 % m/v

CONTENTS:

BATCH NO:

DATE OF MANUFACTURE:

MANUFACTURED AND REGISTERED BY: Nulandis® (A division of AECI Limited)

Reg. No. 1924/002590/06
P.O. Box 14418
Witfield, 1467
Tel : (011) 823 8000
www.nulandis.com



NULANDIS



DIRECTIONS FOR USE: USE ONLY AS DIRECTED

- **Zalexin** should not be added with any sulphur or phosphite based products.

APPLICATION TABLE:

CROP	APPLICATION RATE	APPLICATION METHOD
APPLES, PEARS	200-300 ml/100 ℓ water to a maximum of 3 ℓ/ha	Apply 3-4 applications from 50% flowering and repeat with 21 day intervals. For use on Pears: Do not combine Zalexin with other foliar feeds or adjuvants.
CITRUS	200-300 ml/100 ℓ water to a maximum of 3 ℓ/ha	Apply 3 - 4 applications from 100 % petal drop and repeat with 28 day intervals.
CUCURBITS (PUMPKINS, WATERMELONS, MELONS AND CUCUMBERS)	500 ml/100 ℓ water to a maximum of 4 ℓ/ha	Apply at flowering and repeat with 21 day intervals.
LETTUCE, CAULIFLOWER, CABBAGE, CHINESE CABBAGE	500 ml/100 ℓ water to a maximum of 3 ℓ/ha	Begin applications prior to onset of head formation and repeat at 14 to 21 day intervals.
PEACHES, NECTARINES, PLUMS, APRICOTS, ALMONDS, OLIVES	200-300 ml/100 ℓ water to a maximum of 3 ℓ/ha	Apply 3-4 applications from 75% petal drop and repeat with 21 to 28 day intervals. If weather conditions favour bacterial infections, additional applications can be considered.
POTATOES	500 ml/100 ℓ water to a maximum of 4 ℓ/ha	Apply at tuber initiation and repeat every 21 to 28 days.
TOMATOES, PEPPERS, BEANS, PEAS, ONIONS	500 ml/100 ℓ water to a maximum of 4 ℓ/ha	Begin applications prior to fruit set and repeat at 14 to 21 day intervals.
TOBACCO	500 ml/100 ℓ water to a maximum of 4 ℓ/ha	Apply 2 – 3 applications bi-weekly, starting at 30 cm height

IMPORTANT: BEFORE APPLYING ZALEXIN, CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURAL ADVISER ON THE USE THEREOF.